

Scattering Spaces and a Decomposition of Continuous Spectral Subspace of N -body Quantum Systems

HITOSHI KITADA

Department of Mathematical Sciences, University of Tokyo
Komaba, Meguro, Tokyo 153-8914, Japan
e-mail: kitada@ms.u-tokyo.ac.jp, <http://kims.ms.u-tokyo.ac.jp/>

Dedicated to Professor Yoshimi Saitō on his 60th birthday

December 7, 1999

ABSTRACT. We introduce the notion of scattering space S_b^r for N -body quantum mechanical systems, where b is a cluster decomposition with $2 \leq |b| \leq N$ and r is a real number $0 \leq r \leq 1$. Utilizing these spaces, we give a decomposition of continuous spectral subspace by S_b^1 for N -body quantum systems with long-range pair potentials $V_\alpha^L(x_\alpha) = O(|x_\alpha|^{-\epsilon})$. This is extended to a decomposition by S_b^r with $0 \leq r \leq 1$ for some long-range case. We also prove a characterization of ranges of wave operators by S_b^0 .

Introduction

After Dereziński [D] proved the asymptotic completeness of N -body quantum systems with long-range pair potentials $V_\alpha^L(x_\alpha) = O(|x_\alpha|^{-\epsilon})$ for $\epsilon > \sqrt{3} - 1$, Yafaev [Y] gave an example where the asymptotic completeness breaks down for $0 < \epsilon < 1/2$. It seems that the cause which prevents the asymptotic completeness in his case is some “self-similarity” of approximate threshold eigenvectors by which property the channels corresponding to those eigenvectors cannot be separated well enough as in the usual case. The present paper is partly a reminiscence of my preprint [K3] in 1984, while apart from the problem of asymptotic completeness. After some preparations of section 1, we define in section 2 scattering spaces S_b^r , $2 \leq |b| \leq N$, $0 \leq r \leq 1$, for a given N -body Hamiltonian H . Roughly speaking S_b^r consists of vectors f such that in its evolution $e^{-itH}f$, the distance between clusters in b evolves linearly in t , whereas the size of each cluster in b is bounded by a constant times t^r . We then prove in section 4 a decomposition of the continuous spectral subspace as an orthogonal sum of the scattering spaces S_b^1 , $2 \leq |b| \leq N$. We extend the decomposition to the one by S_b^r with general order $0 \leq r \leq 1$ for some long-range pair potentials by utilizing the results of Dereziński [D]. We also see that the orthogonal projection $E_b(r)$ onto the scattering space S_b^r ($0 \leq r \leq 1$) is discontinuous at some $r = r_0 \in (\epsilon, 1/2)$ in the case of Yafaev’s

potentials [Y]. We prove in section 5 a characterization of the ranges of wave operators: $\mathcal{R}(W_b^\pm) = S_b^0$ for general long-range potentials, where we adopt the wave operators W_b^\pm with time-independent modifiers J_b extending [IK] to N -body case. We assume some decay assumptions on threshold eigenvectors (Assumption 1.2) in addition to the usual assumptions on the decay of pair potentials (Assumption 1.1).

1. Preliminaries

We consider the Schrödinger operator defined in $L^2(R^{\nu N})$ ($\nu \geq 1, N \geq 2$)

$$H = H_0 + V, \quad H_0 = - \sum_{i=1}^N \frac{\hbar^2}{2m_i} \frac{\partial^2}{\partial r_i^2}. \quad (1.1)$$

Here

$$V = \sum_{\alpha} V_{\alpha}(x_{\alpha}), \quad (1.2)$$

where $x_{\alpha} = r_i - r_j$, $r_i = (r_{i1}, \dots, r_{i\nu}) \in R^{\nu}$ is the position vector of the i -th particle, $\frac{\partial}{\partial r_i} = \left(\frac{\partial}{\partial r_{i1}}, \dots, \frac{\partial}{\partial r_{i\nu}} \right)$, $\frac{\partial^2}{\partial r_i^2} = \sum_{j=1}^{\nu} \frac{\partial^2}{\partial r_{ij}^2} = \Delta_{r_i}$, $m_i > 0$ is the mass of the i -th particle, and $\alpha = \{i, j\}$ is a pair with $1 \leq i < j \leq N$. Our assumption on the decay rate of the pair potentials $V_{\alpha}(x_{\alpha})$ is as follows.

Assumption 1.1. $V_{\alpha}(x)$ ($x \in R^{\nu}$) is split into a sum of a real-valued C^{∞} function $V_{\alpha}^L(x)$ and a real-valued measurable function $V_{\alpha}^S(x)$ of $x \in R^{\nu}$ satisfying the following conditions: There are real numbers ϵ and ϵ_1 with $0 < \epsilon, \epsilon_1 < 1$ such that for all multi-indices β

$$|\partial_x^{\beta} V_{\alpha}^L(x)| \leq C_{\beta} \langle x \rangle^{-|\beta| - \epsilon} \quad (1.3)$$

with some constants $C_{\beta} > 0$ independent of $x \in R^{\nu}$, and

$$\langle x \rangle^{1 + \epsilon_1} V_{\alpha}^S(x) (-\Delta_x + 1)^{-1} \text{ is a bounded operator in } L^2(R^{\nu}). \quad (1.4)$$

Here Δ_x is a Laplacian with respect to x , and $\langle x \rangle$ is a C^{∞} function of x such that $\langle x \rangle = |x|$ for $|x| \geq 1$ and $\geq \frac{1}{2}$ for $|x| < 1$.

We can adopt weaker conditions on the differentiability and decay rate for higher derivatives of the long-range part $V_{\alpha}^L(x)$, but for later convenience of exposition, we adopt this form in the present paper.

The free part H_0 of H in (1.1) has various forms in accordance with our choice of coordinate systems. We use the so-called Jacobi coordinates. The center of mass of our N -particle system is

$$X_C = \frac{m_1 r_1 + \dots + m_N r_N}{m_1 + \dots + m_N},$$

and the Jacobi coordinates are defined by

$$x_i = r_{i+1} - \frac{m_1 r_1 + \dots + m_i r_i}{m_1 + \dots + m_i}, \quad i = 1, 2, \dots, N-1. \quad (1.5)$$

Accordingly the corresponding canonically conjugate momentum operators are defined by

$$P_C = \frac{\hbar}{i} \frac{\partial}{\partial X_C}, \quad p_i = \frac{\hbar}{i} \frac{\partial}{\partial x_i}.$$

Using these new X_C, P_C, x_i, p_i , we can rewrite H_0 in (1.1) as

$$H_0 = \tilde{H}_0 + H_C. \quad (1.6)$$

Here

$$\tilde{H}_0 = \sum_{i=1}^{N-1} \frac{1}{2\mu_i} p_i^2 = - \sum_{i=1}^{N-1} \frac{\hbar^2}{2\mu_i} \Delta_{x_i}, \quad H_C = \frac{1}{\sum_{j=1}^N m_j} P_C^2,$$

where $\mu_i > 0$ is the reduced mass defined by the relation:

$$\frac{1}{\mu_i} = \frac{1}{m_{i+1}} + \frac{1}{m_1 + \cdots + m_i}.$$

The new coordinates give a decomposition $L^2(R^{\nu N}) = L^2(R^\nu) \otimes L^2(R^n)$ with $n = \nu(N-1)$ and in this decomposition, H is decomposed

$$H = H_C \otimes I + I \otimes \tilde{H}, \quad \tilde{H} = \tilde{H}_0 + V.$$

H_C is a Laplacian, so we consider \tilde{H} in the Hilbert space $\mathcal{H} = L^2(R^n) = L^2(R^{\nu(N-1)})$. We write this \tilde{H} as H in the followings:

$$H = H_0 + V = \sum_{i=1}^{N-1} \frac{1}{2\mu_i} p_i^2 + \sum_{\alpha} V_{\alpha}(x_{\alpha}) = - \sum_{i=1}^{N-1} \frac{\hbar^2}{2\mu_i} \Delta_{x_i} + \sum_{\alpha} V_{\alpha}(x_{\alpha}). \quad (1.7)$$

This means that we consider the Hamiltonian H in (1.1) restricted to the subspace of $R^{\nu N}$:

$$(m_1 + \cdots + m_N) X_C = m_1 r_1 + \cdots + m_N r_N = 0. \quad (1.8)$$

We equip this subspace with the inner product:

$$\langle x, y \rangle = \sum_{i=1}^{N-1} \mu_i x_i \cdot y_i, \quad (1.9)$$

where \cdot denotes the Euclidean scalar product. With respect to this inner product, the changes of variables between Jacobi coordinates in (1.5) are realized by orthogonal transformations on the space R^n defined by (1.8), while μ_i and x_i depend on the order of the construction of the Jacobi coordinates in (1.5). If we define velocity operator v by

$$v = (v_1, \cdots, v_{N-1}) = (\mu_1^{-1} p_1, \cdots, \mu_{N-1}^{-1} p_{N-1}),$$

we can write using the inner product above

$$H_0 = \frac{1}{2} \langle v, v \rangle. \quad (1.10)$$

Next we introduce clustered Jacobi coordinate. Let $a = \{C_1, \dots, C_k\}$ be a disjoint decomposition of the set $\{1, 2, \dots, N\}$: $C_j \neq \emptyset$ ($j = 1, 2, \dots, k$), $\cup_{j=1}^k C_j = \{1, 2, \dots, N\}$ with $C_i \cap C_j = \emptyset$ when $i \neq j$. We denote the number of elements of a set S by $|S|$. Then $|a| = k$ in the present case, and we call a a cluster decomposition with $|a|$ clusters $C_1, \dots, C_{|a|}$. A clustered Jacobi coordinate $x = (x_a, x^a)$ associated with a cluster decomposition $a = \{C_1, \dots, C_k\}$ is obtained by first choosing a Jacobi coordinate

$$x^{(C_\ell)} = (x_1^{(C_\ell)}, \dots, x_{|C_\ell|-1}^{(C_\ell)}) \in R^{\nu(|C_\ell|-1)} \quad (\ell = 1, 2, \dots, k)$$

for the $|C_\ell|$ particles in the cluster C_ℓ and then by choosing an intercluster Jacobi coordinate

$$x_a = (x_1, \dots, x_{k-1}) \in R^{\nu(k-1)}$$

for the center of mass of the k clusters C_1, \dots, C_k . Then $x^a = (x^{(C_1)}, \dots, x^{(C_k)}) \in R^{\nu(N-k)}$ and $x = (x_a, x^a) \in R^{\nu(N-1)} = R^n$. The corresponding canonically conjugate momentum operator is

$$\begin{aligned} p &= (p_a, p^a), \quad p_a = (p_1, \dots, p_{k-1}), \quad p^a = (p^{(C_1)}, \dots, p^{(C_k)}), \\ p_i &= \frac{\hbar}{i} \frac{\partial}{\partial x_i}, \quad p^{(C_\ell)} = (p_1^{(C_\ell)}, \dots, p_{|C_\ell|-1}^{(C_\ell)}), \quad p_i^{(C_\ell)} = \frac{\hbar}{i} \frac{\partial}{\partial x_i^{(C_\ell)}}. \end{aligned}$$

Accordingly $\mathcal{H} = L^2(R^n)$ is decomposed:

$$\begin{aligned} \mathcal{H} &= \mathcal{H}_a \otimes \mathcal{H}^a, \quad \mathcal{H}_a = L^2(R_{x_a}^{\nu(k-1)}), \\ \mathcal{H}^a &= L^2(R_{x^a}^{\nu(N-k)}) = \mathcal{H}^{(C_1)} \otimes \dots \otimes \mathcal{H}^{(C_k)}, \quad \mathcal{H}^{(C_\ell)} = L^2(R_{x^{(C_\ell)}}^{\nu(|C_\ell|-1)}). \end{aligned} \quad (1.11)$$

In this coordinates system, H_0 in (1.7) is decomposed:

$$\begin{aligned} H_0 &= T_a + H_0^a, \\ T_a &= - \sum_{\ell=1}^{k-1} \frac{\hbar^2}{2M_\ell} \Delta_{x_\ell}, \\ H_0^a &= \sum_{\ell=1}^k H_0^{(C_\ell)}, \quad H_0^{(C_\ell)} = - \sum_{i=1}^{|C_\ell|-1} \frac{\hbar^2}{2\mu_i^{(C_\ell)}} \Delta_{x_i^{(C_\ell)}}, \end{aligned} \quad (1.12)$$

where Δ_{x_ℓ} and $\Delta_{x_i^{(C_\ell)}}$ are ν -dimensional Laplacians and M_ℓ and $\mu_i^{(C_\ell)}$ are the reduced masses. We introduce the inner product in the space $R^n = R^{\nu(N-1)}$ as in (1.9):

$$\begin{aligned} \langle x, y \rangle &= \langle (x_a, x^a), (y_a, y^a) \rangle = \langle x_a, y_a \rangle + \langle x^a, y^a \rangle \\ &= \sum_{\ell=1}^{k-1} M_\ell x_\ell \cdot y_\ell + \sum_{\ell=1}^k \sum_{i=1}^{|C_\ell|-1} \mu_i^{(C_\ell)} x_i^{(C_\ell)} \cdot y_i^{(C_\ell)}, \end{aligned} \quad (1.13)$$

and velocity operator

$$v = (v_a, v^a) = M^{-1}p = (m_a^{-1}p_a, (\mu^a)^{-1}p^a), \quad (1.14)$$

where $M = \begin{pmatrix} m_a & 0 \\ 0 & \mu^a \end{pmatrix}$ is the $n = \nu(N-1)$ dimensional diagonal mass matrix whose diagonals are given by $M_1, \dots, M_{k-1}, \mu_1^{(C_1)}, \dots, \mu_{|C_k|-1}^{(C_k)}$. Then H_0 is written as

$$H_0 = \frac{1}{2}\langle v, v \rangle = T_a + H_0^a = \frac{1}{2}\langle v_a, v_a \rangle + \frac{1}{2}\langle v^a, v^a \rangle. \quad (1.15)$$

We need a notion of order in the set of cluster decompositions. A cluster decomposition b is called a refinement of a cluster decomposition a , iff any $C_\ell \in b$ is a subset of some $D_k \in a$. When b is a refinement of a we denote this as $b \leq a$. $b \not\leq a$ is its negation: some cluster $C_\ell \in b$ is not a subset of any $D_k \in a$. Thus for a pair $\alpha = \{i, j\}$, $\alpha \leq a$ means that $\alpha = \{i, j\} \subset D_k$ for some $D_k \in a$, and $\alpha \not\leq a$ means that $\alpha = \{i, j\} \not\subset D_k$ for any $D_k \in a$. $b < a$ means that $b \leq a$ but $b \neq a$.

We decompose the potential term V in (1.7) as

$$\sum_{\alpha} V_{\alpha}(x_{\alpha}) = V_a + I_a, \quad (1.16)$$

where

$$\begin{aligned} V_a &= \sum_{C_\ell \in a} V_{C_\ell}, \\ V_{C_\ell} &= \sum_{\alpha \subset C_\ell} V_{\alpha}(x_{\alpha}), \\ I_a &= \sum_{\alpha \not\leq a} V_{\alpha}(x_{\alpha}). \end{aligned} \quad (1.17)$$

By definition, V_{C_ℓ} depends only on the variable $x^{(C_\ell)}$ inside the cluster C_ℓ . Similarly, V_a depends only on the variable $x^a = (x^{(C_1)}, \dots, x^{(C_k)}) \in R^{3(N-|a|)}$, while I_a depends on all components of the variable x .

Then H in (1.7) is decomposed:

$$\begin{aligned} H &= H_a + I_a = T_a \otimes I + I \otimes H^a + I_a, \\ H_a &= H - I_a = T_a \otimes I + I \otimes H^a, \\ H^a &= H_0^a + V_a = \sum_{C_\ell \in a} H^{(C_\ell)}, \quad H^{(C_\ell)} = H_0^{(C_\ell)} + V_{C_\ell}, \end{aligned} \quad (1.18)$$

where T_a is an operator in $\mathcal{H}_a = L^2(R_{x_a}^{\nu(k-1)})$, H^a and H_0^a are operators in $\mathcal{H}^a = L^2(R_{x^a}^{\nu(N-k)})$, and $H^{(C_\ell)}$ and $H_0^{(C_\ell)}$ are operators in $\mathcal{H}^{(C_\ell)} = L^2(R_{x^{(C_\ell)}}^{\nu(|C_\ell|-1)})$.

We denote by P_a the orthogonal projection onto the pure point spectral subspace (or eigenspace) $\mathcal{H}_{pp}^a = \mathcal{H}_{pp}(H^a) \subset \mathcal{H}^a$ for H^a . We use the same notation P_a for the

obvious extension $I \otimes P_a$ to the total space \mathcal{H} . For $|a| = N$, we set $P_a = I$. Let $M = 1, 2, \dots$ and P_a^M denote an M -dimensional partial projection of P_a such that $\text{s-lim}_{M \rightarrow \infty} P_a^M = P_a$. We define for $\ell = 1, \dots, N-1$ and an ℓ -dimensional multi-index $M = (M_1, \dots, M_\ell)$ ($M_j \geq 1$)

$$\widehat{P}_\ell^M = \left(I - \sum_{|a_\ell|=\ell} P_{a_\ell}^{M_\ell} \right) \cdots \left(I - \sum_{|a_2|=2} P_{a_2}^{M_2} \right) (I - P^{M_1}). \quad (1.19)$$

(Note that for $|a| = 1$, $a = \{C\}$ with $C = \{1, 2, \dots, N\}$. Thus P^{M_1} is an M_1 -dimensional partial projection into the eigenspace of H .) We further define for a $|a|$ -dimensional multi-index $M_a = (M_1, \dots, M_{|a|-1}, M_{|a|}) = (\widehat{M}_a, M_{|a|})$

$$\widetilde{P}_a^{M_a} = P_a^{M_{|a|}} \widehat{P}_{|a|-1}^{\widehat{M}_a}, \quad 2 \leq |a| \leq N. \quad (1.20)$$

Then it is clear that

$$\sum_{2 \leq |a| \leq N} \widetilde{P}_a^{M_a} = \widehat{P}_1^{M_1} = I - P^{M_1}, \quad (1.21)$$

provided that the component M_j of M_a depends only on the number j but not on a . In the following we use such M_a 's only.

Related with those notions, we denote by $\mathcal{H}_c = \mathcal{H}_c(H)$ the orthogonal complement $\mathcal{H}_{pp}(H)^\perp$ of the eigenspace $\mathcal{H}_{pp} = \mathcal{H}_{pp}(H)$ for the total Hamiltonian H . Namely $\mathcal{H}_c(H)$ is the continuous spectral subspace for H . We note that $\mathcal{H}_c(H) = (I - P_a)\mathcal{H}$ for a unique a with $|a| = 1$, and that for $f \in \mathcal{H}$, $(I - P^{M_1})f \rightarrow (I - P_a)f \in \mathcal{H}_c(H)$ as $M_1 \rightarrow \infty$. We use freely the notations of functional analysis for selfadjoint operators, e.g. $E_H(\Delta)$ is the spectral measure for H .

To state a theorem due to Enss [E2], we introduce an assumption:

Assumption 1.2. For any cluster decomposition a with $2 \leq |a| \leq N-1$ and any integer $M = 1, 2, \dots$,

$$\| |x^a|^2 P_a^M \| < \infty. \quad (1.22)$$

This assumption is concerned with the decay rate of eigenvectors of subsystem Hamiltonians. Since it is known that non-threshold eigenvectors decay exponentially (see Froese and Herbst [FH]), this assumption is the one about threshold eigenvectors.

Let v_a , as above, denote the velocity operator between the clusters in a . It is expressed as $v_a = m_a^{-1} p_a$ for some $\nu(|a|-1)$ -dimensional diagonal mass matrix m_a . Then we can state the theorem.

Theorem 1.3([E2]). *Let $N \geq 2$ and let H be the Hamiltonian H in (1.7) or (1.18) for an N -body quantum-mechanical system. Let Assumptions 1.1 and 1.2 be satisfied. Let $f \in \mathcal{H}$. Then there exist a sequence $t_m \rightarrow \pm\infty$ (as $m \rightarrow \pm\infty$) and a sequence M_a^m of multi-indices whose components all tend to ∞ as $m \rightarrow \pm\infty$ such that for all*

cluster decompositions a with $2 \leq |a| \leq N$, for all $\varphi \in C_0^\infty(R_{x_a}^{\nu(|a|-1)})$, $R > 0$, and $\alpha = \{i, j\} \not\subseteq a$

$$\left\| \frac{|x^a|^2}{t_m^2} \tilde{P}_a^{M_a^m} e^{-it_m H/\hbar} f \right\| \rightarrow 0 \quad (1.23)$$

$$\|F(|x_\alpha| < R) \tilde{P}_a^{M_a^m} e^{-it_m H/\hbar} f\| \rightarrow 0 \quad (1.24)$$

$$\|(\varphi(x_a/t_m) - \varphi(v_a)) \tilde{P}_a^{M_a^m} e^{-it_m H/\hbar} f\| \rightarrow 0 \quad (1.25)$$

as $m \rightarrow \pm\infty$. Here $F(S)$ is the characteristic function of the set defined by the condition S .

We denote the sum of the sets of thresholds and eigenvalues of H by \mathcal{T} :

$$\mathcal{T} = \bigcup_{1 \leq |a| \leq N} \sigma_p(H^a) = \tilde{\mathcal{T}} \cup \sigma_p(H), \quad \tilde{\mathcal{T}} = \bigcup_{2 \leq |a| \leq N} \sigma_p(H^a) \quad (1.26)$$

where

$$\sigma_p(H^a) = \{\tau_1 + \cdots + \tau_{|a|} \mid \tau_\ell \in \sigma_p(H^{(C_\ell)}) \ (C_\ell \in a)\} \quad (1.27)$$

is the set of eigenvalues of a subsystem Hamiltonian $H^a = \sum_{C_\ell \in a} H^{(C_\ell)}$. For $|a| = N$ we define $\sigma_p(H^a) = \{0\}$. Similarly \mathcal{T}_a and $\tilde{\mathcal{T}}_a$ are defined:

$$\mathcal{T}_a = \bigcup_{b \leq a} \sigma_p(H^b) = \tilde{\mathcal{T}}_a \cup \sigma_p(H^a), \quad \tilde{\mathcal{T}}_a = \bigcup_{b < a} \sigma_p(H^b). \quad (1.28)$$

It is known (Froese and Herbst [FH]) that these sets are subsets of $(-\infty, 0]$. Further these sets form bounded, closed and countable subsets of R^1 , and $\sigma_p(H^a)$ accumulates only at $\tilde{\mathcal{T}}_a$ (see Cycon *et al.* [C]).

We use the notation $\Delta \Subset \Delta'$ for Borel sets $\Delta, \Delta' \subset R^k$ to mean that the closure $\bar{\Delta}$ of Δ is compact in R^k and is a subset of the interior of Δ' .

2. Scattering Spaces

In the following we consider the case $t \rightarrow \infty$ only. The other case $t \rightarrow -\infty$ is treated similarly. We also choose a unit system such that $\hbar = 1$. We use the notation $f(t) \sim g(t)$ as $t \rightarrow \infty$ to mean that $\|f(t) - g(t)\| \rightarrow 0$ as $t \rightarrow \infty$ for \mathcal{H} -valued functions $f(t)$ and $g(t)$ of $t > 1$.

Definition 2.1. Let real numbers r, σ, δ and a cluster decomposition b satisfy $0 \leq r \leq 1$, $\sigma, \delta > 0$ and $2 \leq |b| \leq N$.

i) Let $\Delta \Subset R^1 - \mathcal{T}$ be a closed set. We define $S_b^{r\sigma\delta}(\Delta)$ for $0 < r \leq 1$ by

$$S_b^{r\sigma\delta}(\Delta) = \{f \in E_H(\Delta)\mathcal{H} \mid e^{-itH}f \sim \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma t)F(|x^b| \leq \delta t^r)e^{-itH}f \text{ as } t \rightarrow \infty\}. \quad (2.1)$$

For $r = 0$ we define $S_b^{0\sigma}(\Delta)$ by

$$S_b^{0\sigma}(\Delta) = \{f \in E_H(\Delta)\mathcal{H} \mid \lim_{R \rightarrow \infty} \limsup_{t \rightarrow \infty} \left\| e^{-itH}f - \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma t)F(|x^b| \leq R)e^{-itH}f \right\| = 0\}. \quad (2.2)$$

We then define the localized scattering space $S_b^r(\Delta)$ of order $r \in (0, 1]$ for H as the closure of

$$\bigcup_{\sigma > 0} \bigcap_{\delta > 0} S_b^{r\sigma\delta}(\Delta) = \{f \in E_H(\Delta)\mathcal{H} \mid \exists \sigma > 0, \forall \delta > 0 : e^{-itH}f \sim \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma t)F(|x^b| \leq \delta t^r)e^{-itH}f \text{ as } t \rightarrow \infty\}. \quad (2.3)$$

$S_b^0(\Delta)$ is defined as the closure of

$$\bigcup_{\sigma > 0} S_b^{0\sigma}(\Delta) = \{f \in E_H(\Delta)\mathcal{H} \mid \exists \sigma > 0 : \lim_{R \rightarrow \infty} \limsup_{t \rightarrow \infty} \left\| e^{-itH}f - \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma t)F(|x^b| \leq R)e^{-itH}f \right\| = 0\}. \quad (2.4)$$

ii) We define the scattering space S_b^r of order $r \in [0, 1]$ for H as the closure of

$$\bigcup_{\Delta \Subset R^1 - \mathcal{T}} S_b^r(\Delta). \quad (2.5)$$

We note that $S_b^{r\sigma\delta}(\Delta)$, $S_b^{0\sigma}(\Delta)$, $S_b^r(\Delta)$ and S_b^r define closed subspaces of $E_H(\Delta)\mathcal{H}$ and $\mathcal{H}_c(H)$, respectively.

Proposition 2.2. *Let $\Delta \Subset R^1 - \mathcal{T}$ and $f \in S_b^{r\sigma\delta}(\Delta)$ for $0 < r \leq 1$ or $f \in S_b^{0\sigma}(\Delta)$ for $r = 0$ with $\sigma, \delta > 0$ and $2 \leq |b| \leq N$. Then the following limit relations hold:*

i) Let $\alpha \not\leq b$. Then for $0 < r \leq 1$ we have when $t \rightarrow \infty$

$$F(|x_\alpha| < \sigma t)F(|x^b| \leq \delta t^r)e^{-itH}f \rightarrow 0. \quad (2.6)$$

For $r = 0$ we have

$$\lim_{R \rightarrow \infty} \limsup_{t \rightarrow \infty} \left\| F(|x_\alpha| < \sigma t)F(|x^b| \leq R)e^{-itH}f \right\| = 0. \quad (2.7)$$

ii) For $0 < r \leq 1$ we have when $t \rightarrow \infty$

$$F(|x^b| > \delta t^r) e^{-itH} f \rightarrow 0. \quad (2.8)$$

For $r = 0$

$$\lim_{R \rightarrow \infty} \limsup_{t \rightarrow \infty} \|F(|x^b| > R) e^{-itH} f\| = 0. \quad (2.9)$$

iii) There exists a sequence $t_m \rightarrow \infty$ as $m \rightarrow \infty$ depending on $f \in S_b^{r\sigma\delta}(\Delta)$ or $f \in S_b^{0\sigma}(\Delta)$ such that

$$\|(\varphi(x_b/t_m) - \varphi(v_b)) e^{-it_m H} f\| \rightarrow 0 \quad \text{as } m \rightarrow \infty \quad (2.10)$$

for any function $\varphi \in C_0^\infty(R_{x_b}^{\nu(|b|-1)})$.

Proof. i) and ii) are clear from the definition of $S_b^{r\sigma\delta}(\Delta)$ or $S_b^{0\sigma}(\Delta)$. We prove iii). Since $f \in E_H(\Delta)\mathcal{H} \subset H_c(H)$, we have by (1.21), Theorem 1.3 and $f \in S_b^{r\sigma\delta}(\Delta)$ (or $f \in S_b^{0\sigma}(\Delta)$)

$$e^{-it_m H} f \sim \sum_{d \leq b} \tilde{P}_d^{M_d^m} e^{-it_m H} f \quad (2.11)$$

along some sequence $t_m \rightarrow \infty$ depending on f . On each state on the right-hand side (RHS) of (2.11), (1.25) with a replaced by d holds. By the restriction $d \leq b$ in the sum of the RHS of (2.11), we obtain (2.10). \square

The following propositions are obvious by definition.

Proposition 2.3. *Let $2 \leq |b| \leq N$. If $1 \geq r' \geq r > 0$, $\sigma \geq \sigma' > 0$ and $\delta' \geq \delta > 0$ and $\Delta \in R^1 - \mathcal{T}$, then $S_b^{0\sigma}(\Delta) \subset S_b^{0\sigma'}(\Delta)$, $S_b^{0\sigma}(\Delta) \subset S_b^{r\sigma\delta}(\Delta) \subset S_b^{r'\sigma'\delta'}(\Delta)$, $S_b^0(\Delta) \subset S_b^r(\Delta) \subset S_b^{r'}(\Delta)$, $S_b^0(\Delta) \subset S_b^r(\Delta) \subset S_b^r$, and $S_b^0 \subset S_b^r \subset S_b^{r'}$.*

Proposition 2.4. *Let b and b' be different cluster decompositions: $b \neq b'$. Then for any $0 \leq r \leq 1$, S_b^r and $S_{b'}^r$ are orthogonal mutually: $S_b^r \perp S_{b'}^r$.*

3. A Partition of Unity

To state a proposition that will play a fundamental role in our decomposition of continuous spectral subspace by S_b^1 , we prepare some notations. Let b be a cluster decomposition with $2 \leq |b| \leq N$. For any two clusters C_1 and C_2 in b , we define a vector z_{b1} that connects the two centers of mass of the clusters C_1 and C_2 . The number of such vectors when we move over all pairs of clusters in b is $k_b = \binom{|b|}{2}$ in total. We denote these vectors by $z_{b1}, z_{b2}, \dots, z_{bk_b}$.

Let z_{bk} ($1 \leq k \leq k_b$) connect two clusters C_ℓ and C_m in b ($\ell \neq m$). Then for any pair $\alpha = \{i, j\}$ with $i \in C_\ell$ and $j \in C_m$, the vector $x_\alpha = x_{ij}$ is expressed like $(z_{bk}, x^{(C_\ell)}, x^{(C_m)}) \in R^{\nu+\nu(|C_\ell|-1)+\nu(|C_m|-1)}$, where $x^{(C_\ell)} (\in R^{\nu(|C_\ell|-1)})$ and $x^{(C_m)} (\in R^{\nu(|C_m|-1)})$ are the positions of the particles i and j in C_ℓ and C_m , respectively. The vector expression $x_\alpha = (z_{bk}, x^{(C_\ell)}, x^{(C_m)})$ is in the space $R^{\nu+\nu(|C_\ell|-1)+\nu(|C_m|-1)}$. If we

express it in the larger space $R_{z_{bk}}^\nu \times R_{x^b}^{\nu(N-|b|)}$, it would be $x_\alpha = (z_{bk}, x^b)$, and $|x_\alpha|^2 = |z_{bk}|^2 + |x^b|^2$. Thus if $|z_{bk}|^2$ is sufficiently large compared to $|x^b|^2 \geq |x^{(C_\ell)}|^2 + |x^{(C_m)}|^2$, e.g. if $|z_{bk}|^2 > \rho > 0$ and $|x^b|^2 < \theta$ with $\rho \gg \theta > 0$ (which means that ρ/θ is sufficiently large), then $|x_\alpha|^2 > \rho/2$ for all $\alpha \not\leq b$.

Next if $c < b$ and $|c| = |b| + 1$, then just one cluster, say $C_\ell \in b$, is decomposed into two clusters C'_ℓ and C''_ℓ in c , and other clusters in b remain the same in the finer cluster decomposition c . In this case, we can choose just one vector z_{ck} ($1 \leq k \leq k_c$) that connects clusters C'_ℓ and C''_ℓ in c , and we can express $x^b = (z_{ck}, x^c)$. The norm of this vector is written as

$$|x^b|^2 = |z_{ck}|^2 + |x^c|^2. \quad (3.1)$$

Similarly the norm of $x = (x_b, x^b)$ is written as

$$|x|^2 = |x_b|^2 + |x^b|^2. \quad (3.2)$$

We recall that norm is defined, as usual, from the inner product defined by (1.13) which changes in accordance with the cluster decomposition used in each context. E.g., in (3.1), the left-hand side (LHS) is defined by using (1.13) for the cluster decomposition b and the RHS is by using (1.13) for c .

With these preparations, we state the following lemma, which is partially a repetition of [K1, Lemma 2.1]. We define subsets $T_b(\rho, \theta)$ and $\tilde{T}_b(\rho, \theta)$ of $R^n = R^{\nu(N-1)}$ for cluster decompositions b with $2 \leq |b| \leq N$ and real numbers ρ, θ with $1 > \rho, \theta > 0$:

$$T_b(\rho, \theta) = \left(\bigcap_{k=1}^{k_b} \{x \mid |z_{bk}|^2 > \rho|x|^2\} \right) \cap \{x \mid |x_b|^2 > (1 - \theta)|x|^2\}, \quad (3.3)$$

$$\tilde{T}_b(\rho, \theta) = \left(\bigcap_{k=1}^{k_b} \{x \mid |z_{bk}|^2 > \rho\} \right) \cap \{x \mid |x_b|^2 > 1 - \theta\}. \quad (3.4)$$

Subsets S and S_θ ($\theta > 0$) of $R^n = R^{\nu(N-1)}$ are defined by

$$S = \{x \mid |x|^2 \geq 1\},$$

$$S_\theta = \{x \mid 1 + \theta \geq |x|^2 \geq 1\}.$$

Lemma 3.1. *Suppose that constants $1 \geq \theta_1 > \rho_j > \theta_j > \rho_N > 0$ satisfy $\theta_{j-1} \geq \theta_j + \rho_j$ for $j = 2, 3, \dots, N-1$. Then the followings hold:*

i)

$$S \subset \bigcup_{2 \leq |b| \leq N} T_b(\rho_{|b|}, \theta_{|b|}). \quad (3.5)$$

ii) Let $\gamma_j > 1$ ($j = 1, 2$) satisfy

$$\gamma_1 \gamma_2 < r_0 := \min_{2 \leq j \leq N-1} \{\rho_j / \theta_j\}. \quad (3.6)$$

If $b \not\leq c$ with $|b| \geq |c|$, then

$$T_b(\gamma_1^{-1}\rho_{|b|}, \gamma_2\theta_{|b|}) \cap T_c(\gamma_1^{-1}\rho_{|c|}, \gamma_2\theta_{|c|}) = \emptyset. \quad (3.7)$$

iii) For $\gamma > 1$ and $2 \leq |b| \leq N$

$$\begin{aligned} T_b(\rho_{|b|}, \theta_{|b|}) \cap S_{\theta_{N-1}} &\subset \tilde{T}_b(\rho_{|b|}, \theta_{|b|}) \cap S_{\theta_{N-1}} \\ &\subseteq \tilde{T}_b(\gamma^{-1}\rho_{|b|}, \gamma\theta_{|b|}) \cap S_{\theta_{N-1}} \\ &\subset T_b(\gamma_1'^{-1}\rho_{|b|}, \gamma_2'\theta_{|b|}) \cap S_{\theta_{N-1}}, \end{aligned} \quad (3.8)$$

where

$$\gamma_1' = \gamma(1 + \theta_{N-1}), \quad \gamma_2' = (1 + \gamma)(1 + \theta_{N-1})^{-1}. \quad (3.9)$$

iv) If $\frac{2\gamma_1'\gamma_2'}{2-\gamma_1'} < r_0$, then for $2 \leq |b| \leq N$

$$T_b(\gamma_1'^{-1}\rho_{|b|}, \gamma_2'\theta_{|b|}) \subset \{x \mid |x_\alpha|^2 > \rho_{|b|}|x|^2/2 \text{ for all } \alpha \not\leq b\}. \quad (3.10)$$

v) If $\gamma(1 + \gamma) < r_0$ and $b \not\leq c$ with $|b| \geq |c|$, then

$$T_b(\gamma_1'^{-1}\rho_{|b|}, \gamma_2'\theta_{|b|}) \cap T_c(\gamma_1'^{-1}\rho_{|c|}, \gamma_2'\theta_{|c|}) = \emptyset. \quad (3.11)$$

Proof. To prove (3.5), suppose that $|x|^2 \geq 1$ and x does not belong to the set

$$A = \bigcup_{2 \leq |b| \leq N-1} \left[\left(\bigcap_{k=1}^{k_b} \{x \mid |z_{bk}|^2 > \rho_{|b|}|x|^2\} \right) \cap \{x \mid |x_b|^2 > (1 - \theta_{|b|})|x|^2\} \right].$$

Under this assumption, we prove $|x_\alpha|^2 > \rho_N|x|^2$ for all pairs $\alpha = \{i, j\}$. (Note that z_{bk} for $|b| = N$ equals some x_α .) Let $|b| = 2$ and write $x = (z_{b1}, x^b)$. Then by (3.1), $1 \leq |x|^2 = |z_{b1}|^2 + |x^b|^2$. Since x belongs to the complement A^c of the set A , we have $|z_{b1}|^2 \leq \rho_{|b|}|x|^2$ or $|x_b|^2 \leq (1 - \theta_{|b|})|x|^2$. If $|z_{b1}|^2 \leq \rho_{|b|}|x|^2$, then $|x^b|^2 = |x|^2 - |z_{b1}|^2 \geq (1 - \rho_{|b|})|x|^2 \geq (\theta_1 - \rho_{|b|})|x|^2 \geq \theta_{|b|}|x|^2$ by $\theta_{j-1} \geq \theta_j + \rho_j$. Thus $|x_b|^2 = |x|^2 - |x^b|^2 \leq (1 - \theta_{|b|})|x|^2$ for all b with $|b| = 2$.

Next let $|c| = 3$ and assume $|x_c|^2 > (1 - \theta_{|c|})|x|^2$. Then by $x \in A^c$, we can choose z_{ck} with $1 \leq k \leq k_c$ such that $|z_{ck}|^2 \leq \rho_{|c|}|x|^2$. Let C_ℓ and C_m be two clusters in c connected by z_{ck} , and let b be the cluster decomposition obtained by combining C_ℓ and C_m into one cluster with retaining other clusters of c in b . Then $|b| = 2$, $x^b = (z_{ck}, x^c)$, and $|x^b|^2 = |z_{ck}|^2 + |x^c|^2$. Thus $|x_b|^2 = |x|^2 - |x^b|^2 = |x|^2 - |z_{ck}|^2 - |x^c|^2 = |x_c|^2 - |z_{ck}|^2 > (1 - \theta_{|c|} - \rho_{|c|})|x|^2 \geq (1 - \theta_{|b|})|x|^2$, which contradicts the result of the previous step. Thus $|x_c|^2 \leq (1 - \theta_{|c|})|x|^2$ for all c with $|c| = 3$.

Repeating this procedure, we finally arrive at $|x_d|^2 \leq (1 - \theta_{|d|})|x|^2$, thus $|x^d|^2 = |x|^2 - |x_d|^2 \geq \theta_{|d|}|x|^2 > \rho_N|x|^2$ for all d with $|d| = N - 1$. Namely $|x_\alpha|^2 > \rho_N|x|^2$ for all pairs $\alpha = \{i, j\}$. The proof of (3.5) is complete.

We next prove (3.7). By $b \not\leq c$, we can take a pair $\alpha = \{i, j\}$ and clusters $C_\ell, C_m \in c$ such that $\alpha \leq b$, $i \in C_\ell$, $j \in C_m$, and $\ell \neq m$. Then we can write $x_\alpha = (z_{ck}, x^c)$ for some $1 \leq k \leq k_c$. Thus if there is $x \in T_b(\gamma_1^{-1}\rho_{|b|}, \gamma_2\theta_{|b|}) \cap T_c(\gamma_1^{-1}\rho_{|c|}, \gamma_2\theta_{|c|})$, then

$$\gamma_2\theta_{|b|}|x|^2 > |x^b|^2 \geq |x_\alpha|^2 = |z_{bk}|^2 + |x^c|^2 \geq |z_{ck}|^2 > \gamma_1^{-1}\rho_{|c|}|x|^2. \quad (3.12)$$

But since $|b| \geq |c|$, we have $\rho_{|c|} > \gamma_1\gamma_2\theta_{|b|}$ when $|b| = |c|$ by (3.6), and $\rho_{|c|} > \gamma_1\gamma_2\theta_{|c|} \geq \gamma_1\gamma_2(\theta_{|b|} + \rho_{|b|}) > \gamma_1\gamma_2\theta_{|b|}$ when $|c| < |b|$ by $\theta_{j-1} \geq \theta_j + \rho_j$, which both contradict the inequality (3.12). This completes the proof of (3.7).

(3.8) follows by a simple calculation from the inequality $|x|^2(1 + \theta_{N-1})^{-1} \leq 1$ that holds on $S_{\theta_{N-1}}$. (3.10) follows from the relation $|x_\alpha|^2 = |z_{bk}|^2 + |x^b|^2$ stated before the lemma, and (3.11) from $\gamma'_1\gamma'_2 = \gamma(1 + \gamma)$ and ii). \square

In the followings we fix constants $\gamma > 1$ and $1 \geq \theta_1 > \rho_j > \theta_j > \rho_N > 0$ such that

$$\theta_{j-1} \geq \theta_j + \rho_j \quad (j = 2, 3, \dots, N-1), \quad (3.13)$$

$$\max \left\{ \gamma(1 + \gamma), \frac{2\gamma'_1\gamma'_2}{2 - \gamma'_1} \right\} < r_0 = \min_{2 \leq j \leq N-1} \{\rho_j/\theta_j\}, \quad (3.14)$$

where γ'_j ($j = 1, 2$) are defined by (3.9).

Let $\rho(\lambda) \in C^\infty(R^1)$ be such that $0 \leq \rho(\lambda) \leq 1$, $\rho(\lambda) = 1$ ($\lambda \leq -1$), $\rho(\lambda) = 0$ ($\lambda \geq 0$), and $\rho'(\lambda) \leq 0$. Then we define functions $\phi_\sigma(\lambda < \tau)$ and $\phi_\sigma(\lambda > \tau)$ of $\lambda \in R^1$ by

$$\begin{aligned} \phi_\sigma(\lambda < \tau) &= \rho((\lambda - (\tau + \sigma))/\sigma), \\ \phi_\sigma(\lambda > \tau) &= 1 - \phi_\sigma(\lambda < \tau - \sigma) \end{aligned} \quad (3.15)$$

for constants $\sigma > 0, \tau \in R^1$. We note that $\phi_\sigma(\lambda < \tau)$ and $\phi_\sigma(\lambda > \tau)$ satisfy

$$\phi_\sigma(\lambda < \tau) = \begin{cases} 1 & (\lambda \leq \tau) \\ 0 & (\lambda \geq \tau + \sigma) \end{cases} \quad (3.16)$$

$$\phi_\sigma(\lambda > \tau) = \begin{cases} 0 & (\lambda \leq \tau - \sigma) \\ 1 & (\lambda \geq \tau) \end{cases} \quad (3.17)$$

$$\phi'_\sigma(\lambda < \tau) = \frac{d}{d\lambda} \phi_\sigma(\lambda < \tau) \leq 0,$$

$$\phi'_\sigma(\lambda > \tau) \geq 0.$$

We define for a cluster decomposition b with $2 \leq |b| \leq N$

$$\varphi_b(x_b) = \prod_{k=1}^{k_b} \phi_\sigma(|z_{bk}|^2 > \rho_{|b|}) \phi_\sigma(|x_b|^2 > 1 - \theta_{|b|}), \quad (3.18)$$

where $\sigma > 0$ is fixed as

$$0 < \sigma < \min_{2 \leq j \leq N-1} \{(1 - \gamma^{-1})\rho_N, (1 - \gamma^{-1})\rho_j, (\gamma - 1)\theta_j\}. \quad (3.19)$$

Then $\varphi_b(x_b)$ satisfies for $x \in S_{\theta_{N-1}}$

$$\varphi_b(x_b) = \begin{cases} 1 & \text{for } x \in \tilde{T}_b(\rho_{|b|}, \theta_{|b|}), \\ 0 & \text{for } x \notin \tilde{T}_b(\gamma^{-1}\rho_{|b|}, \gamma\theta_{|b|}). \end{cases} \quad (3.20)$$

We set for $|b| = k$ ($k = 2, 3, \dots, N$)

$$J_b(x) = \varphi_b(x_b) \left(1 - \sum_{|b_{k-1}|=k-1} \varphi_{b_{k-1}}(x_{b_{k-1}}) \right) \cdots \left(1 - \sum_{|b_2|=2} \varphi_{b_2}(x_{b_2}) \right). \quad (3.21)$$

By v) and iii) of Lemma 3.1 and (3.20), the sums on the RHS remain only in the case $b < b_j$ for $j = k-1, \dots, 2$ and $x \in S_{\theta_{N-1}}$:

$$J_b(x) = \varphi_b(x_b) \left(1 - \sum_{\substack{|b_{k-1}|=k-1 \\ b < b_{k-1}}} \varphi_{b_{k-1}}(x_{b_{k-1}}) \right) \cdots \left(1 - \sum_{\substack{|b_2|=2 \\ b < b_2}} \varphi_{b_2}(x_{b_2}) \right). \quad (3.22)$$

Thus $J_b(x)$ is a function of the variable x_b only:

$$J_b(x) = J_b(x_b) \quad \text{when } x = (x_b, x^b) \in S_{\theta_{N-1}}. \quad (3.23)$$

We also note that the supports of φ_{b_j} in each sum on the RHS of (3.21) are disjoint mutually in $S_{\theta_{N-1}}$ by iii) and v) of Lemma 3.1. By (3.5) and (3.8) of lemma 3.1, and the definition (3.18)-(3.21) of $J_b(x_b)$, we therefore have

$$\sum_{2 \leq |b| \leq N} J_b(x_b) = 1 \quad \text{on } S_{\theta_{N-1}}.$$

We have constructed a partition of unity on $S_{\theta_{N-1}}$:

Proposition 3.2. *Let real numbers $1 \geq \theta_1 > \rho_j > \theta_j > \rho_N > 0$ satisfy $\theta_{j-1} \geq \theta_j + \rho_j$ for $j = 2, 3, \dots, N-1$. Assume that (3.14) hold and let $J_b(x_b)$ be defined by (3.18)-(3.22). Then we have*

$$\sum_{2 \leq |b| \leq N} J_b(x_b) = 1 \quad \text{on } S_{\theta_{N-1}}. \quad (3.24)$$

$J_b(x_b)$ is a C^∞ function of x_b and satisfies $0 \leq J_b(x_b) \leq 1$. Further on $\text{supp } J_b \cap S_{\theta_{N-1}}$ we have

$$|x_\alpha|^2 > \rho_{|b|} |x|^2 / 2 \quad (3.25)$$

for any pair $\alpha \not\leq b$, and

$$\sup_{x \in \mathbb{R}^n, 2 \leq |b| \leq N} |\nabla_{x_b} J_b(x_b)| < \infty \quad (3.26)$$

for each fixed $\sigma > 0$ in (3.18)-(3.19).

Proof. We have only to see (3.25) and (3.26). But (3.25) is clear by (3.8), (3.10), (3.14), (3.20) and (3.21), and (3.26) follows from (3.15), (3.18) and (3.22). \square

4. A Decomposition of Continuous Spectral Subspace

The following theorem gives a decomposition of $\mathcal{H}_c(H)$ by scattering spaces S_b^1 ($2 \leq |b| \leq N$).

Theorem 4.1. *Let Assumptions 1.1 and 1.2 be satisfied. Then we have*

$$\mathcal{H}_c(H) = \bigoplus_{2 \leq |b| \leq N} S_b^1. \quad (4.1)$$

Proof. Since the set

$$\bigcup_{\Delta \in R^1 - \mathcal{T}} E_H(\Delta) \mathcal{H}$$

is dense in $\mathcal{H}_c(H)$, and S_b^1 ($2 \leq |b| \leq N$) are closed and mutually orthogonal, it suffices to prove that any $\Phi(H)f$ with $\Phi \in C_0^\infty(R^1 - \mathcal{T})$ and $f \in \mathcal{H}$ can be decomposed as a sum of the elements f_b^1 in S_b^1 : $\Phi(H)f = \sum_{2 \leq |b| \leq N} f_b^1$.

We divide the proof into two steps. In the first step I), we prove existence of certain time limits. In the second step II), we prove existence of some ‘‘boundary values’’ of those limits, and conclude the proof of decomposition (4.1).

I) Existence of some time limits:

We decompose $\Phi(H)f$ as a finite sum: $\Phi(H)f = \sum_{j_0}^{\text{finite}} \psi_{j_0}(H)f$, where $\psi_{j_0} \in C_0^\infty(R^1 - \mathcal{T})$. In the step I), we will prove the existence of the limit

$$\lim_{t \rightarrow \infty} \sum_{\ell=1}^L e^{itH} G_{b, \lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b, \lambda_\ell}(t) e^{-itH} \psi_{j_0}(H)f, \quad (4.2)$$

under the assumption that $\text{supp } \psi_{j_0} \subset \tilde{\Delta} \Subset \Delta$ for some intervals $\tilde{\Delta} \Subset \Delta \Subset R^1 - \mathcal{T}$ with $E \in \Delta$ and $\text{diam } \Delta < d(E)$, where $d(E) > 0$ is some small constant depending on $E \in R^1 - \mathcal{T}$ and $\text{diam } S$ denotes the diameter of a set $S \subset R^1$. The relevant factors in (4.2) will be defined in the course of the proof. We will write f for $\psi_{j_0}(H)f$ in the followings.

We take $\psi \in C_0^\infty(R^1)$ such that $\psi(\lambda) = 1$ for $\lambda \in \tilde{\Delta}$ and $\text{supp } \psi \subset \Delta$ for the intervals $\tilde{\Delta} \Subset \Delta$ above. Then $f = \psi(H)f = E_H(\Delta)f \in E_H(\Delta)\mathcal{H} \subset \mathcal{H}_c(H)$ and $e^{-itH}f = \psi(H)e^{-itH}f$. Thus we can use the decomposition (1.21) for the sequences t_m and M_b^m in Theorem 1.3:

$$e^{-it_m H} f = \psi(H)e^{-it_m H} f = \psi(H) \sum_{2 \leq |d| \leq N} \tilde{P}_d^{M_d^m} e^{-it_m H} f. \quad (4.3)$$

By Theorem 1.3-(1.24)

$$\psi(H) \tilde{P}_d^{M_d^m} e^{-it_m H} f \sim \psi(H_d) \tilde{P}_d^{M_d^m} e^{-it_m H} f \quad (4.4)$$

as $m \rightarrow \infty$. Since

$$\tilde{P}_d^{M_d^m} = P_d^{M_{|d|}^m} \widehat{P}_{|d|-1}^{M_d^m}, \quad P_d^{M_{|d|}^m} = \sum_{j=1}^{M_{|d|}^m} P_{d,E_j}, \quad (4.5)$$

where P_{d,E_j} is one dimensional eigenprojection for H^d with eigenvalue E_j , the RHS of (4.4) equals

$$\sum_{j=1}^{M_{|d|}^m} \psi(T_d + E_j) P_{d,E_j} \widehat{P}_{|d|-1}^{M_d^m} e^{-it_m H} f. \quad (4.6)$$

By $\text{supp } \psi \subset \Delta \Subset R^1 - \mathcal{T}$, $E_j \in \mathcal{T}$, and $T_d \geq 0$, we can take constants $\Lambda_d > \lambda_d > 0$ independent of $j = 1, 2, \dots$ such that $\Lambda_d \geq T_d \geq \lambda_d$ if $\psi(T_d + E_j) \neq 0$. Set $\Lambda_0 = \max_d \Lambda_d > \lambda_0 = \min_d \lambda_d > 0$ and

$$\Sigma(E) = \{E - \lambda \mid \lambda \in \mathcal{T}, E \geq \lambda\}. \quad (4.7)$$

Note that we can take $\Lambda_0 > \lambda_0 > 0$ so that

$$\Sigma(E) \Subset (\lambda_0, \Lambda_0) \subset (0, \infty). \quad (4.8)$$

Let $\Psi \in C_0^\infty(R^1)$ satisfy $\Psi(\lambda) = 1$ for $\lambda \in [\lambda_0, \Lambda_0]$ and $\text{supp } \Psi \subset [\lambda_0 - \kappa, \Lambda_0 + \kappa]$ for some small constant $\kappa > 0$ such that the set $[\lambda'_0, \lambda_0] \cup [\Lambda_0, \Lambda'_0]$ is bounded away from $\Sigma(E)$, where $\lambda'_0 = \lambda_0 - 2\kappa > 0$ and $\Lambda'_0 = \Lambda_0 + 2\kappa$. Then the RHS of (4.6) equals for any $m = 1, 2, \dots$

$$\Psi^2(T_d) \psi(H_d) \tilde{P}_d^{M_d^m} e^{-it_m H} f. \quad (4.9)$$

On the other hand, by Theorem 1.3-(1.23) and (1.25), we have

$$\frac{|x^d|^2}{t_m^2} \psi(H_d) \tilde{P}_d^{M_d^m} e^{-it_m H} f \sim 0 \quad (4.10)$$

and

$$\Psi^2(T_d) \psi(H_d) \tilde{P}_d^{M_d^m} e^{-it_m H} f \sim \Psi^2(|x_d|^2 / (2t_m^2)) \psi(H_d) \tilde{P}_d^{M_d^m} e^{-it_m H} f \quad (4.11)$$

as $m \rightarrow \infty$, where to see (4.10) we used (1.23) and $i[H^d, |x^d|^2/t^2] = i[H_0^d, |x^d|^2/t^2] = 2A^d/t^2$ where $A^d = (x^d \cdot p^d + p^d \cdot x^d)/2$, and to see (4.11) the fact that $|x_d|^2/t_m^2$ and H_d commute asymptotically as $m \rightarrow \infty$ by (1.25). Thus by $|x|^2 = |x_d|^2 + |x^d|^2$ we have

$$\Psi^2(T_d) \psi(H_d) \tilde{P}_d^{M_d^m} e^{-it_m H} f \sim \Psi^2(|x|^2 / (2t_m^2)) \psi(H_d) \tilde{P}_d^{M_d^m} e^{-it_m H} f \quad (4.12)$$

as $m \rightarrow \infty$. From (4.3)-(4.4), (4.6), (4.9) and (4.12), we obtain

$$e^{-it_m H} f \sim \Psi^2(|x|^2 / (2t_m^2)) e^{-it_m H} f \quad (4.13)$$

as $m \rightarrow \infty$.

Let constants $\gamma > 1$ and $1 \geq \theta_1 > \rho_j > \theta_j > \rho_N > 0$ be fixed such that

$$\theta_{j-1} \geq \theta_j + \rho_j \quad (j = 2, \dots, N-1), \quad (4.14)$$

$$\max \left\{ \gamma(1 + \gamma), \frac{2\gamma'_1\gamma'_2}{2 - \gamma'_1} \right\} < r_0 = \min_{2 \leq j \leq N-1} \{\rho_j/\theta_j\} \quad (4.15)$$

for γ'_j ($j = 1, 2$) defined by (3.9). Set

$$\lambda''_0 = \lambda'_0 \theta_{N-1} > 0 \quad (4.16)$$

with $\lambda'_0 = \lambda_0 - 2\kappa$ defined above. Let $\tau_0 > 0$ satisfy

$$0 < 16\tau_0 < \lambda''_0 (< \lambda'_0 < \lambda_0). \quad (4.17)$$

We take a finite subset $\{\tilde{\lambda}_\ell\}_{\ell=1}^L$ of \mathcal{T} such that

$$\mathcal{T} \subset \bigcup_{\ell=1}^L (\tilde{\lambda}_\ell - \tau_0, \tilde{\lambda}_\ell + \tau_0). \quad (4.18)$$

Then we can choose real numbers $\lambda_\ell \in R^1$, $\tau_\ell > 0$ ($\ell = 1, 2, \dots, L$) and $\sigma_0 > 0$ such that

$$\begin{aligned} \tau_\ell < \tau_0, \quad \sigma_0 < \tau_0, \quad |\lambda_\ell - \tilde{\lambda}_\ell| < \tau_0, \\ \mathcal{T} \subset \bigcup_{\ell=1}^L (\lambda_\ell - \tau_\ell, \lambda_\ell + \tau_\ell), \quad (\lambda_\ell - \tau_\ell, \lambda_\ell + \tau_\ell) \subset (\tilde{\lambda}_\ell - \tau_0, \tilde{\lambda}_\ell + \tau_0), \\ \text{dist}\{(\lambda_\ell - \tau_\ell, \lambda_\ell + \tau_\ell), (\lambda_k - \tau_k, \lambda_k + \tau_k)\} > 4\sigma_0 (> 0) \quad \text{for any } \ell \neq k. \end{aligned} \quad (4.19)$$

We note that for $\ell = 1, \dots, L$

$$\{\Lambda \mid \tau_\ell \leq |\Lambda - (E - \lambda_\ell)| \leq \tau_\ell + 4\sigma_0\} \cap \Sigma(E) = \emptyset. \quad (4.20)$$

Now let the intervals Δ and $\tilde{\Delta}$ be so small that

$$\text{diam } \tilde{\Delta} < \text{diam } \Delta < \tilde{\tau}_0 := \min_{1 \leq \ell \leq L} \{\sigma_0, \tau_\ell\}. \quad (4.21)$$

Returning to (4.6), we have

$$T_d + E_j \in \text{supp } \psi, \quad (4.22)$$

if $\psi(T_d + E_j) \neq 0$ in (4.6). By $\text{supp } \psi \subset \Delta$, $\text{diam } \Delta < \tilde{\tau}_0$, and $E \in \Delta$, we have from (4.22)

$$-\tilde{\tau}_0 \leq T_d - (E - E_j) \leq \tilde{\tau}_0. \quad (4.23)$$

Thus we have asymptotically on each state in (4.6)

$$-2\tilde{\tau}_0 \leq \frac{|x|^2}{t_m^2} - 2(E - E_j) \leq 2\tilde{\tau}_0. \quad (4.24)$$

By (4.19), $E_j \in \mathcal{T}$ is included in just one set $(\lambda_\ell - \tau_\ell, \lambda_\ell + \tau_\ell)$ for some $\ell = \ell(j)$ with $1 \leq \ell(j) \leq L$. Since $|E_j - \lambda_{\ell(j)}| < \tau_{\ell(j)}$, we have using (4.21)

$$-2\tau_{\ell(j)} - 2\sigma_0 \leq \frac{|x|^2}{t_m^2} - 2(E - \lambda_{\ell(j)}) \leq 2\tau_{\ell(j)} + 2\sigma_0 \quad (4.25)$$

on each state in (4.6). Thus

$$\sum_{\ell=1}^L \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) = 1$$

asymptotically as $m \rightarrow \infty$ on (4.6). Now by the same reasoning that led us to (4.13), we see that (4.3) asymptotically equals as $m \rightarrow \infty$

$$\sum_{\ell=1}^L \sum_{2 \leq |d| \leq N} \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \Psi^2(|x|^2/(2t_m^2)) \tilde{P}_d^{M_d^m} e^{-it_m H} f. \quad (4.26)$$

Since $\phi_{\sigma_0}(|x|^2/t_m^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0) = 1$ on $\text{supp } \phi_{\sigma_0}(|x|^2/t_m^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0)$, (4.26) equals

$$\begin{aligned} & \sum_{\ell=1}^L \sum_{2 \leq |d| \leq N} \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0) \\ & \quad \times \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \Psi^2(|x|^2/(2t_m^2)) \tilde{P}_d^{M_d^m} e^{-it_m H} f. \end{aligned} \quad (4.27)$$

Set

$$B = \langle x \rangle^{-1/2} A \langle x \rangle^{-1/2}, \quad A = \frac{1}{2}(x \cdot p + p \cdot x) = \frac{1}{2}(\langle x, v \rangle + \langle v, x \rangle). \quad (4.28)$$

We note by Theorem 1.3-(1.23) and (1.25) that on the state $\tilde{P}_d^{M_d^m} e^{-it_m H} f$

$$B \sim \sqrt{2T_d} \sim \frac{|x|}{t_m} \quad (4.29)$$

asymptotically as $t_m \rightarrow \infty$. Using this, we replace $\phi_{\sigma_0}(|x|^2/t_m^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0)$ by $\phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0)$ in (4.27). Let $\varphi(\lambda) \in C_0^\infty((\sqrt{2(\lambda_0 - 2\kappa)}, \sqrt{2(\lambda_0 + 2\kappa)}))$, $0 \leq \varphi(\lambda) \leq 1$, and $\varphi(\lambda) = 1$ on $[\sqrt{2(\lambda_0 - \kappa)}, \sqrt{2(\lambda_0 + \kappa)}] (\supset \text{supp}$

$\Psi(\lambda^2/2) \cap (0, \infty)$). We insert a factor $\varphi^2(B)$ into (4.27) and then remove the factor $\Psi^2(|x|^2/(2t_m^2))$ using (4.13):

$$\begin{aligned} & \sum_{\ell=1}^L \sum_{2 \leq |d| \leq N} \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)) < 2\tau_\ell + 4\sigma_0 \\ & \quad \times \phi_{\sigma_0}^2(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \varphi^2(B) \tilde{P}_d^{M_d^m} e^{-it_m H} f. \end{aligned} \quad (4.30)$$

On $\text{supp } \phi_{\sigma_0}(|x|^2/t_m^2 - 2(E - \lambda_\ell)) < 2\tau_\ell + 4\sigma_0$ we have

$$0 < 2(E - \lambda_\ell) - 7\tau_0 \leq \frac{|x|^2}{t_m^2} \leq 2(E - \lambda_\ell) + 7\tau_0. \quad (4.31)$$

Since (4.8), $|\lambda_\ell - \tilde{\lambda}_\ell| < \tau_0$ and (4.17) imply

$$\frac{2(E - \lambda_\ell) + 7\tau_0}{2(E - \lambda_\ell) - 7\tau_0} - 1 = \frac{14\tau_0}{2(E - \lambda_\ell) - 7\tau_0} < \frac{14\lambda_0''/16}{30\lambda_0/16 - \lambda_0''} < \theta_{N-1}, \quad (4.32)$$

we can apply the partition of unity in Proposition 3.2 to the ring defined by (4.31). Then we obtain

$$\begin{aligned} e^{-it_m H} f & \sim \sum_{\ell=1}^L \sum_{2 \leq |b| \leq N} \sum_{2 \leq |d| \leq N} J_b(x_b/(r_\ell t_m)) \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)) < 2\tau_\ell + 4\sigma_0 \\ & \quad \times \phi_{\sigma_0}^2(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \varphi^2(B) \tilde{P}_d^{M_d^m} e^{-it_m H} f, \end{aligned} \quad (4.33)$$

where

$$r_\ell = \sqrt{2(E - \lambda_\ell) - 7\tau_0} > 0 \quad (\ell = 1, \dots, L). \quad (4.34)$$

By the property (3.25), only the terms with $d \leq b$ remain in (4.33):

$$\begin{aligned} e^{-it_m H} f & \sim \sum_{\ell=1}^L \sum_{2 \leq |b| \leq N} \sum_{d \leq b} J_b(x_b/(r_\ell t_m)) \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)) < 2\tau_\ell + 4\sigma_0 \\ & \quad \times \phi_{\sigma_0}^2(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \varphi^2(B) \tilde{P}_d^{M_d^m} e^{-it_m H} f. \end{aligned} \quad (4.35)$$

Using Theorem 1.3-(1.25), we replace x_b/t_m by v_b , and at the same time we introduce a pseudodifferential operator into (4.35):

$$P_b(t) = \phi_\sigma(|x_b/t - v_b|^2 < u) \quad (4.36)$$

with $u > 0$ sufficiently small. Then (4.35) becomes

$$\begin{aligned} e^{-it_m H} f & \sim \sum_{\ell=1}^L \sum_{2 \leq |b| \leq N} \sum_{d \leq b} P_b^2(t_m) J_b(v_b/r_\ell) \phi_{\sigma_0}^2(|x|^2/t_m^2 - 2(E - \lambda_\ell)) < 2\tau_\ell + 4\sigma_0 \\ & \quad \times \phi_{\sigma_0}^2(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \varphi^2(B) \tilde{P}_d^{M_d^m} e^{-it_m H} f. \end{aligned} \quad (4.37)$$

We rearrange the order of the factors on the RHS of (4.37) using that the factors mutually commute asymptotically as $m \rightarrow \infty$ by Theorem 1.3. Setting

$$\begin{aligned} G_{b,\lambda_\ell}(t) &= P_b(t) \phi_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)) < 2\tau_\ell + 4\sigma_0 \\ &\times \phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \varphi(B), \end{aligned} \quad (4.38)$$

we obtain

$$e^{-it_m H} f \sim \sum_{\ell=1}^L \sum_{2 \leq |b| \leq N} \sum_{d \leq b} G_{b,\lambda_\ell}(t_m)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t_m) \tilde{P}_d^{M_d^m} e^{-it_m H} f. \quad (4.39)$$

Now by some calculus of pseudodifferential operators and Theorem 1.3 we note that $P_b(t) J_b(v_b/r_\ell)$ yields a partition of unity $\tilde{J}_b(x_b/(r_\ell t))$ asymptotically as $m \rightarrow \infty$ whose support is close to that of $J_b(x_b/(r_\ell t))$. Then we can recover the terms with $d \not\leq b$, and using (1.21), we remove the sum of $\tilde{P}_d^{M_d^m}$ over $2 \leq |d| \leq N$:

$$e^{-it_m H} f \sim \sum_{\ell=1}^L \sum_{2 \leq |b| \leq N} G_{b,\lambda_\ell}(t_m)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t_m) e^{-it_m H} f. \quad (4.40)$$

We note that on the RHS, the support with respect to $B^2/2$ of the derivative $\phi'_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0)$ is disjoint with $\Sigma(E)$ by (3.16) and (4.20), and the support of $\varphi'(B)$ is similar by (4.8) and the definition of φ above.

We prove the existence of the limit

$$f_{b,\ell} := \lim_{t \rightarrow \infty} e^{itH} G_{b,\lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f \quad (4.41)$$

for $\ell = 1, \dots, L$ and b with $2 \leq |b| \leq N$.

For this purpose we differentiate the function

$$(e^{itH} G_{b,\lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f, g) \quad (4.42)$$

with respect to t , where $f, g \in E_H(\Delta)\mathcal{H}$. Then writing

$$D_t^b g(t) = i[H_b, g(t)] + \frac{dg}{dt}(t) \quad (4.43)$$

for an operator-valued function $g(t)$, we have

$$\begin{aligned}
& \frac{d}{dt} (e^{itH} G_{b,\lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f, g) \\
&= (e^{itH} D_t^b(\varphi(B)) \phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \\
&\quad \times \phi_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0) P_b(t) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f, g) \\
&+ (e^{itH} \varphi(B) D_t^b(\phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0)) \\
&\quad \times \phi_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0) P_b(t) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f, g) \\
&+ (e^{itH} \varphi(B) \phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \\
&\quad \times D_t^b(\phi_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0)) P_b(t) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f, g) \\
&+ (e^{itH} \varphi(B) \phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \\
&\quad \times \phi_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0) D_t^b(P_b(t)) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f, g) \\
&+ ((h.c.)f, g) \\
&+ (e^{itH} i[I_b, G_{b,\lambda_\ell}(t) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t)] e^{-itH} f, g), \tag{4.44}
\end{aligned}$$

where $(h.c.)$ denotes the adjoint of the operator in the terms preceding it.

We need the following lemmas (see [K1, Lemmas 4.1 and 4.2]):

Lemma 4.2. *Let Assumption 1.1 be satisfied. Let $E \in R^1 - \mathcal{T}$. Let $F(s) \in C_0^\infty(R^1)$ satisfy $0 \leq F \leq 1$ and the condition that the support with respect to $s^2/2$ of $F(s)$ is disjoint with $\Sigma(E)$. Then there is a constant $d(E) > 0$ such that for any interval Δ around E with $\text{diam } \Delta < d(E)$, one has*

$$\int_{-\infty}^{\infty} \left\| \frac{1}{\sqrt{\langle x \rangle}} F(B) e^{-itH} E_H(\Delta) f \right\|^2 dt \leq C \|f\|^2 \tag{4.45}$$

for some constant $C > 0$ independent of $f \in \mathcal{H}$.

Lemma 4.3. *For the pseudodifferential operator $P_b(t)$ defined by (4.36) with $u > 0$, there exist norm continuous bounded operators $S(t)$ and $R(t)$ such that*

$$D_t^b P_b(t) = \frac{1}{t} S(t) + R(t) \tag{4.46}$$

and

$$S(t) \geq 0, \quad \|R(t)\| \leq C \langle t \rangle^{-2} \tag{4.47}$$

for some constant $C > 0$ independent of $t \in R^1$.

We switch to a smaller interval Δ if necessary in the followings when we apply Lemma 4.2.

For the first term on the RHS of (4.44) we have

$$D_t^b(\varphi(B)) = \varphi'(B)i[H_b, B] + R_1 \quad (4.48)$$

with

$$\|(H+i)^{-1}\langle x \rangle^{1/2}i[H_b, B]\langle x \rangle^{1/2}(H+i)^{-1}\| < \infty, \quad (4.49)$$

$$\|(H+i)^{-1}\langle x \rangle R_1 \langle x \rangle (H+i)^{-1}\| < \infty. \quad (4.50)$$

(See section 4 of [K1] for a detailed argument yielding the estimates for the remainder terms R_1 here and $S_1(t)$, etc. below.) By the remark after (4.40), the support with respect to $B^2/2$ of $\varphi'(B)$ is disjoint with $\Sigma(E)$. Hence the condition of Lemma 4.2 is satisfied. Thus using (4.49)-(4.50) and rearranging the order of the factors in the first term on the RHS of (4.44) with some integrable errors, we have by Lemma 4.2:

$$\text{the 1st term} = (e^{itH} B_2^{(1)}(t)^* B_1^{(1)}(t) e^{-itH} f, g) + (e^{itH} S_1(t) e^{-itH} f, g), \quad (4.51)$$

where $B_j^{(1)}(t)$ ($j = 1, 2$) and $S_1(t)$ satisfy

$$\int_{-\infty}^{\infty} \|B_j^{(1)}(t) e^{-itH} f\|^2 dt \leq C \|f\|^2, \quad (4.52)$$

$$\|(H+i)^{-1} S_1(t) (H+i)^{-1}\| \leq C t^{-2} \quad (4.53)$$

for some constant $C > 0$ independent of $f \in E_H(\Delta)\mathcal{H}$ and $t \in R^1$.

Similarly by another remark after (4.40) and Lemma 4.2, we have a similar bound for the second term on the RHS of (4.44):

$$\text{the 2nd term} = (e^{itH} B_2^{(2)}(t)^* B_1^{(2)}(t) e^{-itH} f, g) + (e^{itH} S_2(t) e^{-itH} f, g), \quad (4.54)$$

where $B_j^{(2)}(t)$ ($j = 1, 2$) and $S_2(t)$ satisfy

$$\int_{-\infty}^{\infty} \|B_j^{(2)}(t) e^{-itH} f\|^2 dt \leq C \|f\|^2, \quad (4.55)$$

$$\|(H+i)^{-1} S_2(t) (H+i)^{-1}\| \leq C t^{-2} \quad (4.56)$$

for some constant $C > 0$ independent of $f \in E_H(\Delta)\mathcal{H}$ and $t \in R^1$.

For the third term on the RHS of (4.44), we have

$$\begin{aligned} & \varphi(B)\phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \\ & \quad \times D_t^b(\phi_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0)) \\ & = \frac{2}{t}\varphi(B)\phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0) \\ & \quad \times \left(\frac{A}{t} - \frac{|x|^2}{t^2}\right)\phi'_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0) \\ & \quad + S_3(t), \end{aligned} \quad (4.57)$$

where $S_3(t)$ satisfies

$$\|(H+i)^{-1}S_3(t)(H+i)^{-1}\| \leq Ct^{-2}, \quad t > 1. \quad (4.58)$$

On the support of $\phi'_{\sigma_0}(|x|^2/t^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0)$, we have

$$|x|/t \geq \sqrt{2(E - \lambda_\ell) - 2\tau_\ell - 5\sigma_0} > 0$$

by (4.17) and (4.19). Thus there is a large $T > 1$ such that for $t \geq T$ we have $|x| > 1$ and $\langle x \rangle = |x|$, and hence

$$\begin{aligned} 2 \left(\frac{A}{t} - \frac{|x|^2}{t^2} \right) &= \frac{\langle x \rangle}{t} \left(\frac{x}{\langle x \rangle} \cdot D_x - \frac{|x|}{t} \right) + \left(D_x \cdot \frac{x}{\langle x \rangle} - \frac{|x|}{t} \right) \frac{\langle x \rangle}{t} \\ &= 2 \frac{\langle x \rangle}{t} \left(B - \frac{|x|}{t} \right) + tS_4(t) \end{aligned} \quad (4.59)$$

with $\|S_4(t)\| \leq Ct^{-2}$ for $t \geq T$. By (3.16), we have

$$\text{supp } \phi'_{\sigma_0}(|s| < 2\tau_\ell + 4\sigma_0) \subset I_1 \cup I_2$$

with

$$I_1 = [-2\tau_\ell - 5\sigma_0, -2\tau_\ell - 4\sigma_0], \quad I_2 = [2\tau_\ell + 4\sigma_0, 2\tau_\ell + 5\sigma_0], \quad (4.60)$$

and

$$\phi'_{\sigma_0}(|s| < 2\tau_\ell + 4\sigma_0) \geq 0 \quad \text{for } s \in I_1, \quad (4.61)$$

$$\phi'_{\sigma_0}(|s| < 2\tau_\ell + 4\sigma_0) \leq 0 \quad \text{for } s \in I_2. \quad (4.62)$$

Consider the case $|x|^2/t^2 - 2(E - \lambda_\ell) \in I_2$. Then

$$\frac{|x|^2}{t^2} \in [2(E - \lambda_\ell) + 2\tau_\ell + 4\sigma_0, 2(E - \lambda_\ell) + 2\tau_\ell + 5\sigma_0]. \quad (4.63)$$

By the factor $\varphi(B)\phi_{\sigma_0}(|B^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 2\sigma_0)$, we have

$$B^2 \in [2(E - \lambda_\ell) - 2\tau_\ell - 3\sigma_0, 2(E - \lambda_\ell) + 2\tau_\ell + 3\sigma_0] \quad (4.64)$$

and $B \geq \sqrt{2\lambda'_0} > 0$. Thus

$$B - \frac{|x|}{t} \leq 0. \quad (4.65)$$

Therefore by (4.59) and (4.62), (4.57) is positive in this case up to an integrable error. Similarly we see that (4.57) is positive also in the case $|x|^2/t^2 - 2(E - \lambda_\ell) \in I_1$. Rearranging the order of the factors in the third term on the RHS of (4.44) with an integrable error, we see that it has the form

$$\text{the 3rd term} = (e^{itH} A(t)^* A(t) e^{-itH} f, g) + (e^{itH} S_5(t) e^{-itH} f, g) \quad (4.66)$$

with

$$\|(H+i)^{-1}S_5(t)(H+i)^{-1}\| \leq Ct^{-2}. \quad (4.67)$$

The fourth term on the RHS of (4.44) has a similar form by virtue of Lemma 4.3.

The fifth term $((h.c.)f, f)$ is treated similarly to the terms above.

The sixth term on the RHS of (4.44) satisfies

$$|\text{the 6th term}| \leq Ct^{-1-\min\{\epsilon, \epsilon_1\}} \|f\| \|g\|. \quad (4.68)$$

This estimate follows if we note with using (3.25) and some calculus of pseudodifferential operators as stated after (4.39) that the factor $G_{b, \lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b, \lambda_\ell}(t)$ restricts the coordinates in the region: $|x_\alpha|^2 > \rho_{|b|} |x|^2/2$.

Summarizing we have proved that (4.44) is written as

$$\begin{aligned} & \frac{d}{dt} (e^{itH} G_{b, \lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b, \lambda_\ell}(t) e^{-itH} f, g) \\ &= (e^{itH} A(t)^* A(t) e^{-itH} f, g) + \sum_{k=1}^2 (e^{itH} B_2^{(k)}(t)^* B_1^{(k)}(t) e^{-itH} f, g) + (S_6(t) f, g), \end{aligned} \quad (4.69)$$

where with some constant $C > 0$ independent of $t > T$ and $f \in \mathcal{H}$

$$\int_T^\infty \|B_j^{(k)}(t) e^{-itH} E_H(\Delta) f\|^2 \leq C \|f\|^2, \quad (j, k = 1, 2) \quad (4.70)$$

$$\|(H+i)^{-1}S_6(t)(H+i)^{-1}\| \leq Ct^{-1-\min\{\epsilon, \epsilon_1\}}. \quad (4.71)$$

Integrating (4.69) with respect to t on an interval $[T_1, T_2] \subset [T, \infty)$, we obtain

$$\begin{aligned} & (e^{itH} G_{b, \lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b, \lambda_\ell}(t) e^{-itH} f, g) \Big|_{t=T_1}^{T_2} \\ &= \int_{T_1}^{T_2} (A(t) e^{-itH} f, A(t) e^{-itH} g) dt \\ &+ \sum_{k=1}^2 \int_{T_1}^{T_2} (B_1^{(k)}(t) e^{-itH} f, B_2^{(k)}(t) e^{-itH} g) dt + \int_{T_1}^{T_2} (S_6(t) f, g) dt. \end{aligned} \quad (4.72)$$

Hence using (4.70), (4.71) and the uniform boundedness of $G_{b, \lambda_\ell}(t)$ in $t > 1$, we have

$$\int_{T_1}^{T_2} \|A(t) e^{-itH} g\|^2 dt \leq C \|g\|^2 \quad (4.73)$$

for some constant $C > 0$ independent of $T_2 > T_1 \geq T$ and $g \in E_H(\Delta)\mathcal{H}$.

(4.73) and (4.72) with (4.70) and (4.71) then yield that

$$\left| (e^{itH} G_{b, \lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b, \lambda_\ell}(t) e^{-itH} f, g) \Big|_{t=T_1}^{T_2} \right| \leq \delta(T_1) \|f\| \|g\| \quad (4.74)$$

for some $\delta(T_1) > 0$ with $\delta(T_1) \rightarrow 0$ as $T_2 > T_1 \rightarrow \infty$. This means that the limit

$$\tilde{f}_b^1 = \lim_{t \rightarrow \infty} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} f \quad (4.75)$$

exists for any $f \in E_H(\Delta)\mathcal{H}$ and b with $2 \leq |b| \leq N$ if Δ is an interval sufficiently small around $E \in R^1 - \mathcal{T}$: $\text{diam } \Delta < d(E)$. Then the asymptotic decomposition (4.40) implies

$$f = \sum_{2 \leq |b| \leq N} \tilde{f}_b^1 \quad (4.76)$$

for $f = \psi(H)f = E_H(\Delta)f$. Further by the existence of the limit (4.75) and $f = E_H(\Delta)f$, we see that \tilde{f}_b^1 satisfies

$$E_H(\Delta)\tilde{f}_b^1 = \tilde{f}_b^1 \quad (4.77)$$

in a way similar to the proof of the intertwining property of wave operators.

Now returning to the first $\Phi(H)f$, and noting that $\text{supp } \Phi$ is compact in $R^1 - \mathcal{T}$, we take a finite number of open intervals $\Delta_{j_0} \Subset R^1 - \mathcal{T}$ such that $E_{j_0} \in \Delta_{j_0}$, $\text{diam } \Delta_{j_0} < d(E_{j_0})$, and $\text{supp } \Phi \Subset \bigcup_{j_0}^{\text{finite}} \Delta_{j_0} \Subset R^1 - \mathcal{T}$. Then we can take $\psi_{j_0} \in C_0^\infty(\Delta_{j_0})$ such that $\Phi(H)f = \sum_{j_0}^{\text{finite}} \psi_{j_0}(H)f$. Thus from (4.75)-(4.77), we obtain the existence of the limit for $2 \leq |b| \leq N$:

$$\tilde{f}_b^1 = \lim_{t \rightarrow \infty} \sum_{j_0}^{\text{finite}} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} \psi_{j_0}(H)f, \quad (4.78)$$

and the relations

$$\Phi(H)f = \sum_{2 \leq |b| \leq N} \tilde{f}_b^1, \quad E_H(\Delta)\tilde{f}_b^1 = \tilde{f}_b^1 \quad (4.79)$$

for any set $\Delta \Subset R^1 - \mathcal{T}$ with $\text{supp } \Phi \subset \Delta$.

Set

$$\sigma_j = \sqrt{\gamma^{-1}\rho_j\lambda'_0/2}, \quad \delta_j = \sqrt{\gamma\theta_j\Lambda'_0} \quad (j = 2, 3, \dots, N, \quad \theta_N = 0). \quad (4.80)$$

Then by (4.78), some calculus of pseudodifferential operators, and

$$\text{supp } (J_b(x_b/r_\ell)\phi_{\sigma_0}(|x|^2 - 2(E - \lambda_\ell)| < 2\tau_\ell + 4\sigma_0)) \Subset \tilde{T}_b(\gamma^{-1}\rho_{|b|}, \gamma\theta_{|b|}), \quad (4.81)$$

which follows from (3.19)-(3.21), we see that as $t \rightarrow \infty$

$$\begin{aligned} e^{-itH} \tilde{f}_b^1 &\sim \sum_{k=1}^K \sum_{\ell=1}^L G_{b,\lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} E_H(\Delta_k) f \\ &\sim \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma_{|b|} t) F(|x^b| \leq \delta_{|b|} t) \\ &\quad \times \sum_{k=1}^K \sum_{\ell=1}^L G_{b,\lambda_\ell}(t)^* J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} E_H(\Delta_k) f \\ &\sim \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma_{|b|} t) F(|x^b| \leq \delta_{|b|} t) e^{-itH} \tilde{f}_b^1. \end{aligned} \quad (4.82)$$

(4.79) and (4.82) imply

$$\tilde{f}_b^1 \in S_b^{1\sigma|b|\delta|b|}(\Delta) \quad (2 \leq |b| \leq N). \quad (4.83)$$

II) A refinement:

As in (4.18)-(4.19), we take a finite subset $\{\tilde{\lambda}_\ell^b\}_{\ell=1}^{L_b}$ of \mathcal{T}_b for a constant $\tau_0^b > 0$ with $\tau_0^b < \tau_0$ such that

$$\mathcal{T}_b \subset \bigcup_{\ell=1}^{L_b} (\tilde{\lambda}_\ell^b - \tau_0^b, \tilde{\lambda}_\ell^b + \tau_0^b), \quad (4.84)$$

and choose real numbers $\lambda_\ell^b \in R^1$, $\tau_\ell^b > 0$ ($\ell = 1, \dots, L_b$) and $\sigma_0^b > 0$ such that

$$\begin{aligned} \tau_\ell^b < \tau_0^b, \quad \sigma_0^b < \tau_0^b, \quad |\lambda_\ell^b - \tilde{\lambda}_\ell^b| < \tau_0^b, \\ \mathcal{T}_b \subset \bigcup_{\ell=1}^{L_b} (\lambda_\ell^b - \tau_\ell^b, \lambda_\ell^b + \tau_\ell^b), \quad (\lambda_\ell^b - \tau_\ell^b, \lambda_\ell^b + \tau_\ell^b) \subset (\tilde{\lambda}_\ell^b - \tau_0^b, \tilde{\lambda}_\ell^b + \tau_0^b), \\ \text{dist}\{(\lambda_\ell^b - \tau_\ell^b, \lambda_\ell^b + \tau_\ell^b), (\lambda_k^b - \tau_k^b, \lambda_k^b + \tau_k^b)\} > 4\sigma_0^b (> 0) \quad \text{for any } \ell \neq k. \end{aligned} \quad (4.85)$$

We set $\mathcal{T}_b^F = \{\lambda_\ell\}_{\ell=1}^{L_b}$ and

$$\tilde{\tau}_0^b = \min_{1 \leq \ell \leq L_b} \{\sigma_0^b, \tau_\ell^b\}. \quad (4.86)$$

Then, we take $\psi_1(\lambda) \in C_0^\infty(R^1)$ such that

$$0 \leq \psi_1 \leq 1, \quad (4.87)$$

$$\psi_1(\lambda) = \begin{cases} 1 & \text{for any } \lambda \text{ with } |\lambda - \lambda_\ell| \leq \tilde{\tau}_0^b/2 \text{ for some } \lambda_\ell \in \mathcal{T}_b^F \\ 0 & \text{for any } \lambda \text{ with } |\lambda - \lambda_\ell| \geq \tilde{\tau}_0^b \text{ for all } \lambda_\ell \in \mathcal{T}_b^F \end{cases} \quad (4.88)$$

and we divide (4.78) as follows:

$$\tilde{f}_b^1 = h_b + g_b, \quad (4.89)$$

where

$$h_b = \lim_{t \rightarrow \infty} \sum_{j_0}^{\text{finite}} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* \psi_1(H^b) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} \psi_{j_0}(H) f, \quad (4.90)$$

$$g_b = \lim_{t \rightarrow \infty} \sum_{j_0}^{\text{finite}} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* (I - \psi_1)(H^b) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} \psi_{j_0}(H) f. \quad (4.91)$$

The proof of the existence of these limits is similar to that of \tilde{f}_b^1 in (4.78), since the change in the present case is the appearance of the commutator $[H, \psi_1(H^b)] = [I_b, \psi(H^b)]$ whose treatment is quite the same as that of the commutators including

I_b in (4.68). We introduce the decomposition (1.21) into h_b and g_b on the left of $e^{-itH}\psi_{j_0}(H)f$ as in (4.3). Then by the factor $J_b(v_b/r_\ell)G_{b,\lambda_\ell}(t)$, we see that only the terms with $d \leq b$ in the sum in (4.3) remain asymptotically as $t = t_m \rightarrow \infty$ by the arguments similar to step I). On each summand $P_{d,E_j}\widehat{P}_{|d|-1}^{\widehat{M}_d^m}$ in these terms (see (4.5)), H^b asymptotically equals $H_d^b = T_d^b + H^d = T_d^b + E_j \sim |x_d^b|^2/(2t_m^2) + E_j \sim |x^b|^2/(2t_m^2) + E_j$, where for $d \leq b$, $H_d^b = T_d^b + H^d = H_d - T_b$, $T_d^b = T_d - T_b$ and $x^b = (x_d^b, x^d)$ is a clustered Jacobi coordinate inside the coordinate x^b . Thus we have

$$\begin{aligned} & \psi_1(H^b)J_b(v_b/r_\ell)G_{b,\lambda_\ell}(t)P_{d,E_j}\widehat{P}_{|d|-1}^{\widehat{M}_d^m}e^{-it_m H}\psi_{j_0}(H)f \\ & \sim \psi_1(|x^b|^2/(2t_m^2) + E_j)J_b(v_b/r_\ell)G_{b,\lambda_\ell}(t)P_{d,E_j}\widehat{P}_{|d|-1}^{\widehat{M}_d^m}e^{-it_m H}\psi_{j_0}(H)f \end{aligned} \quad (4.92)$$

as $m \rightarrow \infty$. If $\psi_1(|x^b|^2/(2t_m^2) + E_j) \neq 0$, then for some $\ell = 1, \dots, L_b$

$$\left| \frac{|x^b|^2}{2t_m^2} - (\lambda_\ell^b - E_j) \right| \leq \tilde{\tau}_0^b. \quad (4.93)$$

If ℓ is a (unique) $\ell(j)$ such that $E_j \in (\lambda_{\ell(j)}^b - \tau_{\ell(j)}^b, \lambda_{\ell(j)}^b + \tau_{\ell(j)}^b)$, we have

$$\frac{|x^b|^2}{2t_m^2} \leq \tau_{\ell(j)}^b + \tilde{\tau}_0^b < \tau_0^b + \tilde{\tau}_0^b. \quad (4.94)$$

Thus setting $\delta' = \sqrt{2(\tau_0^b + \tilde{\tau}_0^b)}$, we have

$$|x^b| \leq \delta' t_m. \quad (4.95)$$

If $\ell \neq \ell(j)$, we have by (4.93)

$$0 \leq \lambda_\ell^b - E_j + \tilde{\tau}_0^b,$$

from which and (4.85)-(4.86) follows

$$\lambda_\ell^b - E_j \geq 4\sigma_0^b.$$

Thus from (4.93)

$$\frac{|x^b|^2}{2t_m^2} \geq 4\sigma_0^b - \tilde{\tau}_0^b \geq 3\sigma_0^b \geq 3\tilde{\tau}_0^b. \quad (4.96)$$

Setting $\sigma' = \sqrt{6\tilde{\tau}_0^b}$ we then have for $\ell \neq \ell(j)$

$$|x^b| \geq \sigma' t_m. \quad (4.97)$$

Therefore h_b can be decomposed as

$$h_b = f_b^{\delta'} + g_{b1}^{\sigma'}, \quad (4.98)$$

where

$$f_b^{\delta'} = \lim_{t \rightarrow \infty} \sum_{j_0}^{\text{finite}} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* F(|x^b| \leq \delta't) \psi_1(H^b) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} \psi_{j_0}(H) f, \quad (4.99)$$

$$g_b^{\sigma'} = \lim_{t \rightarrow \infty} \sum_{j_0}^{\text{finite}} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* F(|x^b| \geq \sigma't) \psi_1(H^b) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} \psi_{j_0}(H) f, \quad (4.100)$$

The existence of the limit (4.99) is proved similarly to that of (4.90) by rewriting the factor $F(|x^b| \leq \delta't)$ as a smooth one and absorbing it into $J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t)$ with changing the constants in it suitably. The existence of (4.100) then follows from this, (4.78) and (4.98).

For g_b , similarly to $g_b^{\sigma'}$ we obtain

$$g_b = \lim_{t \rightarrow \infty} \sum_{j_0}^{\text{finite}} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* F(|x^b| \geq \sigma't) (I - \psi_1)(H^b) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} \psi_{j_0}(H) f. \quad (4.101)$$

Setting

$$\begin{aligned} g_b^{\sigma'} &= g_b^{\sigma'} + g_b \\ &= \lim_{t \rightarrow \infty} \sum_{j_0}^{\text{finite}} \sum_{\ell=1}^L e^{itH} G_{b,\lambda_\ell}(t)^* F(|x^b| \geq \sigma't) J_b(v_b/r_\ell) G_{b,\lambda_\ell}(t) e^{-itH} \psi_{j_0}(H) f, \end{aligned} \quad (4.102)$$

we obtain a decomposition of \tilde{f}_b^1 :

$$\tilde{f}_b^1 = f_b^{\delta'} + g_b^{\sigma'}, \quad (4.103)$$

where $f_b^{\delta'}$ and $g_b^{\sigma'}$ satisfy

$$e^{-itH} f_b^{\delta'} \sim \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma_{|b|} t) F(|x^b| \leq \delta't) e^{-itH} f_b^{\delta'}, \quad (4.104)$$

$$e^{-itH} g_b^{\sigma'} \sim \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma_{|b|} t) F(|x^b| \leq \delta_{|b|} t) F(|x^b| \geq \sigma't) e^{-itH} g_b^{\sigma'}. \quad (4.105)$$

We can prove the existence of the limits

$$f_b^1 = \lim_{\delta' \downarrow 0} f_b^{\delta'}, \quad g_b^1 = \lim_{\sigma' \downarrow 0} g_b^{\sigma'} \quad (4.106)$$

in the same way as in Enss [E3, Lemma 4.8], because we can take ψ_1 in (4.87)-(4.88) monotonically decreasing when $\tilde{\tau}_0^b \downarrow 0$ and the factors $F(|x^b| \leq \delta't)$ and $F(|x^b| \geq \sigma't)$

can be treated similarly to ψ_1 by regarding x^b/t as a single variable. Further we have as in (4.77)

$$E_H(\Delta)f_b^1 = f_b^1, \quad E_H(\Delta)g_b^1 = g_b^1, \quad (4.107)$$

which, (4.104) and (4.106) imply

$$f_b^1 \in S_b^1. \quad (4.108)$$

Thus we have a decomposition:

$$\tilde{f}_b^1 = f_b^1 + g_b^1, \quad f_b^1 \in S_b^1. \quad (4.109)$$

$g_b^{\sigma'}$ can be decomposed further by using the partition of unity of the ring $\sigma' \leq |x^b|/t \leq \delta_{|b|}$ with regarding x^b as a total variable x in Proposition 3.2. Arguing similarly to steps I) and II), we can prove that g_b^1 can be decomposed as a sum of the elements f_d^1 of S_d^1 with $d < b$. Combining this with (4.76), (4.108) and (4.109), we obtain (4.1). \square

We remark that Theorem 4.1 implies the asymptotic completeness when the long-range part V_α^L vanishes for all pairs α , because in this case we see straightforwardly that $S_b^1 = \mathcal{R}(W_b^\pm)$, where W_b^\pm are the short-range wave operators defined by

$$W_b^\pm = s\text{-}\lim_{t \rightarrow \pm\infty} e^{itH} e^{-itH_b} P_b. \quad (4.110)$$

For the case when long-range part does not vanish, we have the following

Theorem 4.2. *Let Assumptions 1.1 and 1.2 be satisfied. Then*

i) For $2(2 + \epsilon)^{-1} < r \leq 1$

$$S_b^r = S_b^1. \quad (4.111)$$

ii) If $\epsilon > 2(2 + \epsilon)^{-1}$, i.e. when $\epsilon > \sqrt{3} - 1$, we have for all r with $0 \leq r \leq 1$

$$S_b^r = S_b^1. \quad (4.112)$$

iii) If $\epsilon > 1/2$ and $V_\alpha^L(x_\alpha) \geq 0$ for all pairs α , then we have for all r with $0 \leq r \leq 1$

$$S_b^r = S_b^1. \quad (4.113)$$

Proof. i) and ii) follow from Proposition 5.8 of [D] and Proposition 2.3 above. (4.112) for $r = 0$ follows from the proof of Proposition 5.8 of [D]. iii) follows from Theorem 1.1 and Proposition 4.3 of [K2] and (4.116) below: Note that $\mathcal{R}(\Omega_b^\psi)$ in Theorem 1.1-(1.31) of [K2] constitutes a dense subset of S_b^1 , when ψ varies in $C_0^\infty(R^1 - \mathcal{T})$. \square

From Theorem 4.2-ii), iii) and Theorem 4.1 follows

Theorem 4.3. *Let Assumptions 1.1 and 1.2 be satisfied with $\epsilon > 2(2 + \epsilon)^{-1}$ or with $\epsilon > 1/2$ and $V_\alpha^L(x_\alpha) \geq 0$ for all pairs α . Then we have for all r with $0 \leq r \leq 1$*

$$\bigoplus_{2 \leq |b| \leq N} S_b^r = \mathcal{H}_c(H). \quad (4.114)$$

In the next section, we will construct modified wave operators:

$$W_b^\pm = s\text{-}\lim_{t \rightarrow \pm\infty} e^{itH} J_b e^{-itH_b} P_b \quad (4.115)$$

with J_b being an extension of J of [IK] to the N -body case. We will then prove

$$\mathcal{R}(W_b^\pm) = S_b^0, \quad (4.116)$$

which and Theorems 4.2 and 4.3 imply

Theorem 4.4. *Let Assumptions 1.1 and 1.2 be satisfied with $\epsilon > 2(2 + \epsilon)^{-1}$ or with $\epsilon > 1/2$ and $V_\alpha^L(x_\alpha) \geq 0$ for all pairs α . Then we have for all r with $0 \leq r \leq 1$*

$$\mathcal{R}(W_b^\pm) = S_b^r, \quad (4.117)$$

and

$$\bigoplus_{2 \leq |b| \leq N} \mathcal{R}(W_b^\pm) = \mathcal{H}_c(H). \quad (4.118)$$

One might expect that (4.114) and (4.118) are always true, but it is denied:

Theorem 4.5. *Let Assumptions 1.1 and 1.2 be satisfied and let $N \geq 3$. Then the followings hold:*

i) *Let $2 \leq |b| \leq N$ and let $E_b(r)$ be the orthogonal projection onto S_b^r ($0 \leq r \leq 1$). Then $E_b(r_1) \leq E_b(r_2)$ for $0 \leq r_1 \leq r_2 \leq 1$, and the discontinuous points of $E_b(r)$ with respect to $r \in [0, 1]$ in the strong operator topology are at most countable.*

ii) *Let $0 < \epsilon < 1/2$ in Assumption 1.1. Then there are long-range pair potentials $V_\alpha(x_\alpha)$ such that for some cluster decomposition b with $2 \leq |b| \leq N$, $E_b(r)$ is discontinuous at $r = r_0$, where $\epsilon < r_0 := (\epsilon + 1)/3 < 1/2$. In particular, there are real numbers r_1 and r_2 with $0 \leq r_1 < r_0 < r_2 \leq 1$ such that*

$$S_b^{r_1} \subsetneq S_b^{r_2}. \quad (4.119)$$

Proof. i) By Proposition 2.3, S_b^r ($0 \leq r \leq 1$) is a family of closed subspaces of a separable Hilbert space \mathcal{H} that increases when $r \in [0, 1]$ increases. Thus the corresponding orthogonal projection $E_b(r)$ ($0 \leq r \leq 1$) onto S_b^r increases as r increases, and hence has at most a countable number of discontinuous points with respect to $r \in [0, 1]$ in the strong operator topology.

ii) holds by Theorem 4.3 of [Y], Theorem 4.1 and Proposition 2.3, for b , $|b| = N - 1$, with a suitable choice of pair potentials that satisfy Assumption 1.1. In fact, the sum of the ranges $\mathcal{R}(W_n)$ of Yafaev's wave operators W_n ($n = 1, 2, \dots$) in Theorem 4.3 of [Y] constitutes a subspace of $(E_b(r_0 + 0) - E_b(r_0 - 0))\mathcal{H}$ for b with $|b| = N - 1$ by his construction of W_n , which means that $E_b(r)$ is discontinuous at $r = r_0$. Here $E_b(r_0 \pm 0) = s\text{-}\lim_{r \rightarrow r_0 \pm 0} E_b(r)$. \square

5. A Characterization of the Ranges of Wave Operators

The purpose in this section is to prove relation (4.116) for general long-range pair potentials $V_\alpha(x_\alpha)$ under Assumptions 1.1 and 1.2. The inclusion

$$\mathcal{R}(W_b^\pm) \subset S_b^0 \quad (5.1)$$

is a trivial relation for any form of definition of the wave operators W_b^\pm . Thus our main concern is to prove the reverse inclusion

$$S_b^0 \subset \mathcal{R}(W_b^\pm). \quad (5.2)$$

The proof of this inclusion is essentially the same for any definition of wave operators and is not difficult in the light of Enss method [E1]. As announced, we here consider the wave operators of the form

$$W_b^\pm = s\text{-}\lim_{t \rightarrow \pm\infty} e^{itH} J_b e^{-itH_b} P_b, \quad (5.3)$$

where J_b is an extension of the identification operator or stationary modifier introduced in [IK] for two-body long-range case. The first task in this section is to construct J_b . Our relation (5.2) then follows from the definition of the scattering spaces S_b^0 and properties of J_b by Enss method.

To make the descriptions simple we hereafter consider the case $V_\alpha^S = 0$ for all pairs α . The recovery of the short-range potentials in the following arguments is easy.

Let a C^∞ function $\chi_0(x)$ of $x \in R^\nu$ satisfy

$$\chi_0(x) = \begin{cases} 1 & (|x| \geq 2) \\ 0 & (|x| \leq 1). \end{cases} \quad (5.4)$$

To define J_b we introduce time-dependent potentials $I_{b\rho}(x_b, t)$ for $\rho \in (0, 1)$:

$$I_{b\rho}(x_b, t) = I_b(x_b, 0) \prod_{k=1}^{k_b} \chi_0(\rho z_{bk}) \chi_0(\langle \log \langle t \rangle \rangle z_{bk} / \langle t \rangle). \quad (5.5)$$

Then $I_{b\rho}(x_b, t)$ satisfies

$$|\partial_{x_b}^\beta I_{b\rho}(x_b, t)| \leq C_\beta \rho^{\epsilon_0} \langle t \rangle^{-\ell} \quad (5.6)$$

for any $\ell \geq 0$ and $0 < \epsilon_0 < \epsilon$ with $\epsilon_0 + \ell < |\beta| + \epsilon$, where $C_\beta > 0$ is a constant independent of t, x_b and ρ .

Then we can apply almost the same arguments as in section 2 of [IK] to get a solution $\varphi_b(x_b, \xi_b)$ of the eikonal equation:

$$\frac{1}{2} |\nabla_{x_b} \varphi_b(x_b, \xi_b)|^2 + I_b(x_b, 0) = \frac{1}{2} |\xi_b|^2 \quad (5.7)$$

in some conic region in phase space. More exactly we have the following theorems. Let

$$\cos(z_{bk}, \zeta_{bk}) := \frac{z_{bk} \cdot \zeta_{bk}}{|z_{bk}|_e |\zeta_{bk}|_e},$$

where $|z_{bk}|_e = (z_{bk} \cdot z_{bk})^{1/2}$ is the Euclidean norm. We then set for $R_0, d > 0$ and $\theta \in (0, 1)$

$$\Gamma_\pm(R_0, d, \theta) = \{(x_b, \xi_b) \mid |z_{kb}| \geq R_0, |\zeta_{bk}| \geq d, \pm \cos(z_{bk}, \zeta_{bk}) \geq \theta \ (k = 1, \dots, k_b)\},$$

where ζ_{bk} is the variable conjugate to z_{bk} .

Theorem 5.1. *Let Assumption 1.1 be satisfied with $V_\alpha^S = 0$ for all pairs α . Then there exists a C^∞ function $\phi_b^\pm(x_b, \xi_b)$ that satisfies the following properties: For any $0 < \theta, d < 1$, there exists a constant $R_0 > 1$ such that for any $(x_b, \xi_b) \in \Gamma_\pm(R_0, d, \theta)$*

$$\frac{1}{2}|\nabla_{x_b} \phi_b^\pm(x_b, \xi_b)|^2 + I_b(x_b, 0) = \frac{1}{2}|\xi_b|^2 \quad (5.8)$$

and

$$|\partial_{x_b}^\alpha \partial_{\xi_b}^\beta (\phi_b^\pm(x_b, \xi_b) - x_b \cdot \xi_b)| \leq \begin{cases} C_{\alpha\beta} (\max_{1 \leq k \leq k_b} \langle z_{bk} \rangle)^{1-\epsilon}, & \alpha = 0 \\ C_{\alpha\beta} (\min_{1 \leq k \leq k_b} \langle z_{bk} \rangle)^{1-\epsilon-|\alpha|}, & \alpha \neq 0, \end{cases} \quad (5.9)$$

where $C_{\alpha\beta} > 0$ is a constant independent of $(x_b, \xi_b) \in \Gamma_\pm(R_0, d, \theta)$.

From this we can derive the following theorem in quite the same way as that for Theorem 2.5 of [IK]. Let $0 < \theta < 1$ and let $\psi_\pm(\tau) \in C^\infty([-1, 1])$ satisfy

$$\begin{aligned} 0 &\leq \psi_\pm(\tau) \leq 1, \\ \psi_+(\tau) &= \begin{cases} 1 & \text{for } \theta \leq \tau \leq 1, \\ 0 & \text{for } -1 \leq \tau \leq \theta/2, \end{cases} \\ \psi_-(\tau) &= \begin{cases} 0 & \text{for } -\theta/2 \leq \tau \leq 1, \\ 1 & \text{for } -1 \leq \tau \leq -\theta. \end{cases} \end{aligned}$$

We set

$$\chi_\pm(x_b, \xi_b) = \prod_{k=1}^{k_b} \psi_\pm(\cos(z_{bk}, \zeta_{bk}))$$

and define $\varphi_b(x_b, \xi_b) = \varphi_{b,\theta,d,R_0}(x_b, \xi_b)$ by

$$\begin{aligned} \varphi_b(x_b, \xi_b) &= \{(\phi_b^+(x_b, \xi_b) - x_b \cdot \xi_b)\chi_+(x_b, \xi_b) + (\phi_b^-(x_b, \xi_b) - x_b \cdot \xi_b)\chi_-(x_b, \xi_b)\} \\ &\quad \times \prod_{k=1}^{k_b} \chi_0(2\zeta_{bk}/d)\chi_0(2z_{bk}/R_0) + x_b \cdot \xi_b \end{aligned}$$

for $d, R_0 > 0$. Note that $\varphi_{b,\theta,d,R_0}(x_b, \xi_b) = \varphi_{b,\theta,d',R'_0}(x_b, \xi_b)$ when $|z_{bk}| \geq \max(R_0, R'_0)$, $|\zeta_{bk}| \geq \max(d, d')$ for all k . We then have

Theorem 5.2. *Let Assumption 1.1 be satisfied with $V_\alpha^S = 0$ for all pairs α . Let $0 < \theta < 1$ and $d > 0$. Then there exists a constant $R_0 > 1$ such that the C^∞ function $\varphi_b(x_b, \xi_b)$ defined above satisfies the following properties.*

i) For $(x_b, \xi_b) \in \Gamma_+(R_0, d, \theta) \cup \Gamma_-(R_0, d, \theta)$, φ_b is a solution of

$$\frac{1}{2}|\nabla_{x_b} \varphi_b(x_b, \xi_b)|^2 + I_b(x_b, 0) = \frac{1}{2}|\xi_b|^2. \quad (5.10)$$

ii) For any $(x_b, \xi_b) \in R^{2\nu(|b|-1)}$ and multi-indices α, β , φ_b satisfies

$$|\partial_{x_b}^\alpha \partial_{\xi_b}^\beta (\varphi_b(x_b, \xi_b) - x_b \cdot \xi_b)| \leq \begin{cases} C_{\alpha\beta} (\max\langle z_{bk} \rangle)^{1-\epsilon}, & \alpha = 0 \\ C_{\alpha\beta} (\min\langle z_{bk} \rangle)^{1-\epsilon-|\alpha|}, & \alpha \neq 0. \end{cases} \quad (5.11)$$

In particular, if $\alpha \neq 0$,

$$|\partial_{x_b}^\alpha \partial_{\xi_b}^\beta (\varphi_b(x_b, \xi_b) - x_b \cdot \xi_b)| \leq C_{\alpha\beta} R_0^{-\epsilon_0} (\min\langle z_{bk} \rangle)^{1-\epsilon_1-|\alpha|} \quad (5.12)$$

for any $\epsilon_0, \epsilon_1 \geq 0$ with $\epsilon_0 + \epsilon_1 = \epsilon$. Further

$$\varphi_b(x_b, \xi_b) = x_b \cdot \xi_b \quad \text{when } |z_{bk}| \leq R_0/2 \text{ or } |\zeta_{bk}| \leq d/2 \text{ for some } k. \quad (5.13)$$

iii) Let

$$a_b(x_b, \xi_b) = e^{-i\varphi_b(x_b, \xi_b)} \left(T_b + I_b(x_b, 0) - \frac{1}{2} |\xi_b|^2 \right) e^{i\varphi_b(x_b, \xi_b)}. \quad (5.14)$$

Then

$$a_b(x_b, \xi_b) = \frac{1}{2} |\nabla_{x_b} \varphi_b(x_b, \xi_b)|^2 + I_b(x_b, 0) - \frac{1}{2} |\xi_b|^2 + i(T_b \varphi_b)(x_b, \xi_b) \quad (5.15)$$

and

$$|\partial_{x_b}^\alpha \partial_{\xi_b}^\beta a_b(x_b, \xi_b)| \leq \begin{cases} C_{\alpha\beta} (\min\langle z_{bk} \rangle)^{-1-\epsilon-|\alpha|}, & (x_b, \xi_b) \in \Gamma_+(R_0, d, \theta) \cup \Gamma_-(R_0, d, \theta) \\ C_{\alpha\beta} (\min\langle z_{bk} \rangle)^{-\epsilon-|\alpha|} \langle \xi_b \rangle, & \text{otherwise.} \end{cases} \quad (5.16)$$

We now define $J_b = J_{b, \theta, d, R_0}$ by

$$J_b f(x_b) = (2\pi)^{-\nu(|b|-1)} \int_{R^{\nu(|b|-1)}} \int_{R^{\nu(|b|-1)}} e^{i(\varphi_b(x_b, \xi_b) - y_b \cdot \xi_b)} f(y_b) dy_b d\xi_b \quad (5.17)$$

for $f \in \mathcal{H}_b = L^2(R^{\nu(|b|-1)})$ as an oscillatory integral (see e.g. [KK]). Wave operators W_b^\pm are now defined by

$$W_b^\pm = \text{s-} \lim_{t \rightarrow \pm\infty} e^{itH} J_b e^{-itH_b} P_b. \quad (5.18)$$

We note that this definition depends on θ, d, R_0 , but applying stationary phase method to e^{-itT_b} in $e^{-itH_b} = e^{-itT_b} \otimes e^{-itH^b}$ on the RHS we see that the dependence disappears in the limit $t \rightarrow \pm\infty$ by the remark made just before Theorem 5.2. Further the asymptotic behavior seen by the stationary phase method tells that the inclusion (5.1) holds:

$$\mathcal{R}(W_b^\pm) \subset S_b^0, \quad (5.19)$$

if the limits (5.18) exist. The existence of (5.18) follows from Theorem 5.2-iii), the asymptotic behavior of e^{-itT_b} and Assumptions 1.1-1.2 by noting the relations

$$\begin{aligned} (HJ_b - J_bH_b)e^{-itH_b}P_b f(x) &= ((T_b + I_b(x_b, x^b))J_b - J_bT_b)e^{-itH_b}P_b f(x), \\ ((T_b + I_b(x_b, 0))J_b - J_bT_b)g(x_b) \\ &= (2\pi)^{-\nu(|b|-1)} \int_{R^{\nu(|b|-1)}} \int_{R^{\nu(|b|-1)}} e^{i(\varphi_b(x_b, \xi_b) - y_b \cdot \xi_b)} a_b(x_b, \xi_b) g(y_b) dy_b d\xi_b \end{aligned} \quad (5.20)$$

and the fact that $s\text{-}\lim_{M \rightarrow \infty} P_b^M = P_b$. Thus to prove the reverse inclusion (5.2)

$$S_b^0 \subset \mathcal{R}(W_b^\pm), \quad (5.21)$$

it suffices to prove that

$$f \in S_b^0 \ominus \mathcal{R}(W_b^\pm) \quad (5.22)$$

implies

$$f = 0. \quad (5.23)$$

To see this we consider the case $t \rightarrow +\infty$ and the quantity

$$(I - e^{isH} J_b e^{-isH_b} J_b^{-1}) e^{-itH} f = (J_b - e^{isH} J_b e^{-isH_b}) J_b^{-1} e^{-itH} f \quad (5.24)$$

for $f \in S_b^0(\Delta)$, $\Delta \in R^1 - \mathcal{T}$, and $t, s \geq 0$ and use Enss method. Here the existence of J_b^{-1} follows from Theorem 3.3 of [K4] by taking $R_0 > 0$ in (5.12) large enough (with a slight adaptation to the present case for phases and symbols satisfying the estimates in Theorem 5.2). (5.24) equals

$$-i \int_0^s e^{iuH} (HJ_b - J_bH_b) e^{-iuH_b} J_b^{-1} du e^{-itH} f. \quad (5.25)$$

By Definition 2.1-i)-(2.4) of $S_b^0(\Delta)$, we approximate f by $h \in S_b^{0\sigma}(\Delta)$ for some small $\sigma > 0$ with an arbitrarily small error $\delta > 0$ so that $\|f - h\| < \delta$. Then we have for any sufficiently large $R > 0$

$$\limsup_{t \rightarrow \infty} \left\| e^{-itH} h - \prod_{\alpha \not\leq b} F(|x_\alpha| \geq \sigma t) F(|x^b| \leq R) e^{-itH} h \right\| < \delta. \quad (5.26)$$

Proposition 2.2-iii) and $h \in S_b^{0\sigma}(\Delta)$ yield that for some sequence $t_m \rightarrow \infty$ (as $m \rightarrow \infty$)

$$\|(\varphi(x_b/t_m) - \varphi(v_b))e^{-it_m H} h\| \rightarrow 0 \quad \text{as } m \rightarrow \infty$$

for any $\varphi \in C_0^\infty(R^{\nu(|b|-1)})$. Replacing t and f in (5.25) by t_m and h , we can therefore insert or remove the factor

$$\Phi = \prod_{k=1}^{k_b} Q_k \tilde{F}(|p_{bk}| \geq \sigma') \tilde{F}(|p_b| \leq S) \tilde{F}(|z_{bk}|/t \geq \sigma') \tilde{F}(|x^b| \leq R) \quad (5.27)$$

to or from the left of $e^{-it_m H} h$ in (5.25) anytime with an error $\delta > 0$. Here $p_{bk} = \frac{1}{i} \frac{\partial}{\partial z_{bk}}$, $\sigma' > 0$ is a small number with $\sigma' < \sigma$, $\tilde{F}(|p_b| \leq S)$ comes from $E_H(\Delta)$ in $h = E_H(\Delta)h$, and $\tilde{F}(\tau \leq S)$ is a smooth characteristic function of the set $\{\tau \in R^1 \mid \tau \leq S\}$ with a slope independent of S , and Q_k is a pseudodifferential operator

$$Q_k g(x_b) = (2\pi)^{-\nu(|b|-1)} \int_{R^{\nu(|b|-1)}} \int_{R^{\nu(|b|-1)}} e^{(x_b \cdot \xi_b - y_b \cdot \zeta_b)} q_k(z_{bk}, \zeta_{bk}) g(y_b) dy_b d\zeta_b \quad (5.28)$$

with symbol $q_k(z_{bk}, \zeta_{bk})$ satisfying

$$\begin{aligned} |\partial_{z_{bk}}^\beta \partial_{\zeta_{bk}}^\gamma q_k(z_{bk}, \zeta_{bk})| &\leq C_{\beta\gamma} \langle z_{bk} \rangle^{-|\beta|} \langle \zeta_{bk} \rangle^{-|\gamma|}, \\ q_k(z_{bk}, \zeta_{bk}) &= 0 \quad \text{for} \quad \cos(z_{bk}, \zeta_{bk}) \leq \theta \text{ or } |z_{bk}| \leq R_0. \end{aligned} \quad (5.29)$$

The order of products in (5.25) of factors in (5.27) and J_b^{-1} may be arbitrary because these factors are mutually commutative asymptotically as $t \rightarrow \infty$ by virtue of (5.26). We note that $d > 0$ in the definition of $J_b = J_{b, \theta, d, R_0}$ can be taken smaller than $\sigma' > 0$ beforehand since W_b^+ is independent of $d > 0$ as mentioned. Thus we can assume the following in addition to (5.29):

$$q(z_{bk}, \zeta_{bk}) = 0 \quad \text{for} \quad |\zeta_{bk}| \leq d. \quad (5.30)$$

We now insert the decomposition (1.21) to the left of $e^{-it_m H} h$ in (5.25) with noting $(I - P^{M_1^m})f = f$ by $f \in \mathcal{H}_c(H)$. Then by $\|(I - P^{M_1^m})h - h\| < 2\delta$ and by inserting the factor (5.27) to the left of $e^{-it_m H} h$ after the insertion of (1.21), we have

$$\limsup_{m \rightarrow \infty} \|(I - P_b^{M_{|b|}^m}) \Phi e^{-it_m H} h\| < 3\delta. \quad (5.31)$$

By the factor $\tilde{F}(|x^b| \leq R)$ in (5.27) and $E_H(\Delta)$ in $h = E_H(\Delta)h$, $P_b^{M_{|b|}^m}$ in (5.31) converges to P_b as $m \rightarrow \infty$ in operator norm in the expression (5.31). It thus suffices to consider the quantity

$$\int_0^s e^{iuH} ((T_b + I_b(x_b, x^b))J_b - J_b T_b) e^{-iuH_b} J_b^{-1} du P_b^{M_{|b|}^{m_0}} \Phi e^{-it_m H} h \quad (5.32)$$

for some large but fixed m_0 with an error $\delta > 0$. Since $P_b^{M_{|b|}^{m_0}} = \sum_{j=1}^{M_{|b|}^{m_0}} P_{b, E_j}$ ($0 \leq M_{|b|}^{m_0} < \infty$) with P_{b, E_j} being one dimensional eigenprojection of H^b corresponding to eigenvalue E_j , (5.32) is reduced to considering

$$\int_0^s e^{-iuE_j} e^{iuH} ((T_b + I_b(x_b, x^b))J_b - J_b T_b) P_{b, E_j} e^{-iuT_b} J_b^{-1} \Phi du e^{-it_m H} h. \quad (5.33)$$

By Assumptions 1.1-1.2, the factor P_{b, E_j} bounds the variable x^b and yields a short-range error of order $O((\min \langle z_{bk} \rangle)^{-1-\epsilon})$ on the left of e^{-iuT_b} when we replace $I_b(x_b, x^b)$ by $I_b(x_b, 0)$, and we have that (5.33) equals

$$\begin{aligned} \int_0^s e^{-iuE_j} e^{iuH} P_{b, E_j} O(\langle x^b \rangle) ((T_b + I_b(x_b, 0))J_b - J_b T_b + O((\min \langle z_{bk} \rangle)^{-1-\epsilon})) \\ \times e^{-iuT_b} J_b^{-1} \Phi du e^{-it_m H} h, \end{aligned} \quad (5.34)$$

where $O(\langle x^b \rangle)$ is an operator such that $\langle x^b \rangle^{-1} O(\langle x^b \rangle)$ is bounded. Using (5.20) and the estimate (5.16) in Theorem 5.2-iii) and applying the propagation estimates in Lemma 3.3-ii) of [IK] (again with a slight adaptation to the present case), we now get the estimate:

$$\|((T_b + I_b(x_b, 0))J_b - J_b T_b + O((\min\langle z_{bk} \rangle)^{-1-\epsilon}))e^{-iuT_b} J_b^{-1} \Phi(\min\langle z_{bk} \rangle)^{\epsilon/2}\| \leq C\langle u \rangle^{-1-\epsilon/2} \quad (5.35)$$

for some constant $C > 0$ independent of $u \geq 0$. On the other hand (5.26) yields that

$$\|(\min\langle z_{bk} \rangle)^{-\epsilon/2} e^{-it_m H} h\|$$

is asymptotically less than 2δ as $m \rightarrow \infty$. This and (5.35) prove that the norm of (5.32) is asymptotically less than a constant times δ as $m \rightarrow \infty$.

Returning to (5.24) we have proved that

$$\begin{aligned} & \limsup_{m \rightarrow \infty} \sup_{s \geq 0} \|(I - e^{isH} J_b e^{-isH_b} J_b^{-1})e^{-it_m H} f\| \\ & \approx_\delta \limsup_{m \rightarrow \infty} \sup_{s \geq 0} \|(I - e^{isH} J_b e^{-isH_b} J_b^{-1})P_b^{M_{|b|}^m} e^{-it_m H} f\| \leq C\delta, \end{aligned} \quad (5.36)$$

where $a \approx_\delta b$ means that $|a - b| \leq C\delta$ for some constant $C > 0$. Since wave operator $W_b^+ = \text{s-lim}_{s \rightarrow \infty} e^{isH} J_b e^{-isH_b} P_b$ exists, (5.36) yields

$$\limsup_{m \rightarrow \infty} \|(I - W_b^+ J_b^{-1})P_b^{M_{|b|}^m} e^{-it_m H} f\| \leq C\delta. \quad (5.37)$$

By the arguments above deriving (5.31) we can remove $P_b^{M_{|b|}^m}$ and get

$$\limsup_{m \rightarrow \infty} \|(I - W_b^+ J_b^{-1})e^{-it_m H} f\| \leq C\delta. \quad (5.38)$$

Since we assumed (5.22), f is orthogonal to $\mathcal{R}(W_b^+)$. Thus taking the inner product of the vector inside the norm in (5.38) with $e^{-it_m H} f$, we have

$$\|f\|^2 = \lim_{m \rightarrow \infty} |(e^{-it_m H} f, e^{-it_m H} f)| = \lim_{m \rightarrow \infty} |(e^{-it_m H} f, (I - W_b^+ J_b^{-1})e^{-it_m H} f)| \leq C\delta\|f\|.$$

As $\delta > 0$ is arbitrary, this gives $f = 0$, proving (5.23). The proof of (4.116) is complete.

REFERENCES

- [C] H. L. Cycon *et al.*, *Schrödinger Operators*, Springer-Verlag, 1987.
- [D] J. Dereziński, *Asymptotic completeness of long-range N -body quantum systems*, *Annals of Math.* **138** (1993), 427-476.
- [E1] V. Enss, *Asymptotic completeness for quantum mechanical potential scattering I*, *Commun. Math. Phys.* **61** (1978), 285-291.
- [E2] V. Enss, *Introduction to asymptotic observables for multiparticle quantum scattering*, *Schrödinger Operators*, Aarhus 1985 (E. Balslev, ed.), *Lect. Note in Math.*, vol. 1218, Springer-Verlag, 1986, pp. 61-92.

- [E3] V. Enss, *Long-range scattering of two- and three-body quantum systems*, Equations aux dérivées partielles, Publ. Ecole Polytechnique, Palaiseau (1989), 1-31.
- [FH] R. Froese and I. Herbst, *Exponential bounds and absence of positive eigenvalues for N -body Schrödinger operators*, Commun. Math. Phys. **87** (1982), 429-447.
- [IK] H. Isozaki and H. Kitada, *Modified wave operators with time-independent modifiers*, Journal of the Fac. Sci, University of Tokyo **Sec. IA**, **32** (1985), 77-104.
- [K1] H. Kitada, *Asymptotic completeness of N -body wave operators I. Short-range quantum systems*, Rev. in Math. Phys. **3** (1991), 101-124.
- [K2] H. Kitada, *Asymptotic completeness of N -body wave operators II. A new proof for the short-range case and the asymptotic clustering for long-range systems*, Functional Analysis and Related Topics, 1991 (H. Komatsu, ed.), Lect. Note in Math, vol. 1540, Springer-Verlag, 1993, pp. 149-189.
- [K3] H. Kitada, *Asymptotic completeness for N -body Schrödinger operators I. Short-range potentials*, preprint (1984, February).
- [K4] H. Kitada, *Fourier integral operators with weighted symbols and micro-local resolvent estimates*, J. Math. Soc. Japan **39** (1987), 101-124.
- [KK] H. Kitada and H. Kumano-go, *A family of Fourier integral operators and the fundamental solution for a Schrödinger equation*, Osaka J. Math. **18** (1981), 291-360.
- [Y] D. Yafaev, *New channels in three-body long-range scattering*, Equations aux dérivées partielles Publ. Ecole Polytechnique, Palaiseau (1994), XIV-1—XIV-11.